

headaches, insomnia, and emotional lability. Psychologic testing was normal. A 21-year-old man experienced nondisabling irritability, insomnia, and headaches. Psychiatric and psychologic interviews yielded inconsistent results. The diagnosis was posttraumatic stress disorder.

Nerve damage has been reported in exposed appendages. Fleck (1983) summarized findings of nerve damage in the hands and forearms of a woman who was using a microwave oven that apparently malfunctioned. According to the report, at the end of a cooking cycle she reached into the cavity to retrieve a dish and was purportedly exposed to the output of a 600-W oven for 5 seconds. The nerve damage persisted resulting in pain and heat sensitization of nerve fibers. Apparently no evaluation was made of the microwave oven. Dickason and Barutt (1984) also observed nerve damage, persistent pain, and hypersensitivity to light touch in a woman overexposed while cooking with a microwave oven. In this case, the oven was evaluated and found to be malfunctioning. Exposure lasted 60 to 90 seconds. Ciano et al. (1981) reported paresthesia in the hand of a cafeteria waitress exposed to a malfunctioning microwave oven over the span of 3 days. Tintinalli, Krause, and Gursel (1983) also noted paresthesia in the hand of a man who received a momentary exposure to a malfunctioning microwave oven. Thirty minutes postexposure, the hand was nonresponsive, pale, and cold. Sixty minutes after exposure, the hand responded normally.

Hocking, Joyner, and Fleming (1988) evaluated an incident where a nine-man radio lineman party was exposed to 4.139-GHz microwaves while disconnecting a waveguide from a metal gantry. Estimates of the power density for the two members who experienced the highest exposures were 4.6 mW/cm² at 2 m for about 90 minutes. The estimated skin SAR was 3.8 W/kg, while the whole-body average SAR was 0.13 W/kg. The other men were exposed at <0.15 mW/cm². Claims of behavioral effects, temporary sexual impairment, insomnia, irritability, and loosening of scalp hair were made by the high-exposure group. Behavioral effects were also claimed by

members of the low-exposure group. The authors conclude that medical "examinations have found various, slight (but inconsistent) abnormalities in the staff which could not be related to the high or low level of exposure. It is concluded that this exposure of up to 4.6 mW/cm² at 4.1 GHz for nearly ninety minutes has not resulted in harmful effects."

3.5.8.3 Reproductive Effects

Rosenthal and Beering (1968) reported a case of male infertility. This was attributed to microwave exposure because the individual worked for four years as a maintenance man on energized weather radar systems. He reported sensing the feeling of warmth, but it is not possible to place this in the proper context because no exposure data were provided.

Rubin and Erdman (1959) summarized four cases of the clinical application of 2.45-GHz microwave diathermy to treat pelvic inflammation. Two patients were treated during pregnancy. One of these spontaneously aborted but subsequently became pregnant and delivered a healthy baby. The other patient had a normal baby. The other two were treated prior to pregnancy, and both delivered normal infants.

A microwave oven repairman had plastic induration of the penis, which the investigators note had not been associated with MW exposure. However, they could not "discount a true casual relationship to microwaves." No degenerative testicular changes were observed. Abdominal exposure (2.45 GHz) was reported to be 18 mW/cm² (Rose et al. 1969).

3.5.8.4 Skin Effects

Ciano et al. (1981) observed necrosis of the skin in a young woman who operated a plastic welding machine. Her hand had been caught in an energized sealer for approximately 2 seconds. Damage was extensive, and the hand nonresponsive, which resulted in amputation.

In the case report by Lim et al. (1993), mention is made of facial erythema in a main-

tenance technician. However, no exposure data are given that would positively link this to microwave exposure. As mentioned earlier, the actual power density emitted from the dish antenna is not known, although estimates from normal operation would be low. Computer modeling of a worst-case leak during normal operation (30 mW/cm²) would produce a small temperature rise (0.48°C) that "could not realistically cause the reported injury" (Hocking, Joyner, Fleming, and Anderson 1994). At a lower frequency, 2.45 GHz, Rose and colleagues (1969) have reported a "thermally-induced erythema *ab igne*" in microwave-oven repairmen with facial exposure around 20 mW/cm², although the duration of exposure is not clear (*ab igne* means the erythema is localized and due to heat). Gellin (1971) terms this facial radiodermatitis, which includes erythema and "a sense of heat."

There was no report of a sensation of heat or pain in the account by Lim et al (1993), although other reports of erythema in the literature commonly include such descriptions. In the report by Fleck (1983) of exposure to the hands and arms due to a malfunctioning (600 W) consumer microwave oven, the subject perceived the exposure as follows: "She felt a hot pulsating sensation and burning in all fingers and finger nails, as well as a sensation of 'needles' from forearms to fingers." Subsequently she experienced erythema, edema, and "jabbing pain." In another case where a microwave oven malfunctioned, the subject "experienced a sudden burning pain of the radial side of her hand" and later developed redness and swelling (Dickason and Barutt 1984).

Brodin and Bleiberg (1973) reported two cases of cutaneous damage associated with microwave oven use. The women, who worked at a snack bar, exhibited deformity and ridging of the fingernails. Both women operated the same oven and the onset of the damage occurred at the same time. It is not clear how long the women were exposed, but there appear to have been multiple instances of exposure over a number of months. During the purported exposure, there was no sensation of heat or pain or report of erythema or edema,

which would be expected. It was alleged that the oven was malfunctioning, but an evaluation could not be performed, because it had been returned to the manufacturer. Following this, the "lesions began to improve when the unit was removed."

Hocking, Joyner, and Fleming (1988) report loosening of scalp hair in a group of radiolinenmen, but this was not linked to RF exposure.

A number of occurrences of painful skin damage associated with spark discharge have been investigated by federal OSHA inspectors (Hagaman 1989a; Shepich 1990; Heins 1990). Tissue damage occurs when RF energy is rapidly released just prior to skin contact with a conductive object in which the incident RF field has induced currents. One occurrence was when construction workers were building a highway in Hawaii beneath a coast-guard navigation antenna. "Radio frequency signals transmitted from the antenna may induce a perceptible electrical charge in large rubber tire vehicles, construction equipment and ungrounded metallic structures" (Hagaman 1989a). An energy threshold of more than 15 μ J was believed to be necessary to produce the effect (Hagaman 1989b). Similar problems were also reported for longshoremen who contacted RF-energized parts of cranes and cables. In these reports the RF sources were AM radio station transmitting towers (Shepich 1990; Heins 1990; Heins and Curtis 1990).

3.5.8.5 Death

A case of death related to microwave exposure near a radar transmitter was reported by McLaughlin (1957). The individual reported extreme tenderness and pain in the abdominal cavity. Upon admission to the hospital, the patient received an appendectomy but died ten days after the operation. Merckel (1972) and Ely (1971, 1985) have disputed McLaughlin's claim that this death was associated with overexposure to MW radiation. Merckel (1972) attributes it to appendicitis. Ely (1985) notes that at the operational frequency (10 GHz), there would have been con-

siderable skin damage due to the shallow penetration depth, and this was not observed by McLaughlin.

3.5.8.6 Cancer

Archimbaud and colleagues (1989) in Lyon, France, reported a single case of acute myelogenous leukemia in a 46-year-old man who had repaired 3-kW microwave oven generators for a number of years. Thirty to thirty-six generators were repaired weekly, and part of the process required that power be supplied to each generator was for about 1 minute. The patient estimated that he was exposed about 5 min/d to microwaves from an unprotected generator. No other exposure information (location, distance, power density, etc.) was provided. In conclusion, Archimbaud et al. state, "Intense exposure to microwave radiations should therefore be considered as a potential aetiology of secondary leukaemia."

This conclusion has been criticized by Jauchem (1990) for lack of exposure information and how the literature cited by Archimbaud et al. (1989) in support of their conclusion was interpreted. In a response, Archimbaud (1990) explained that during exposure, the patient "was standing on the side of the generator while testing it by heating a glass of water" and that "he did not recall any abnormal heat sensation," which would be expected with high-level exposure at this frequency. In a critique, Hocking and Garson (1990) note that deep burns and cataracts would also be expected with intense exposure.

A concern has been raised dealing with the potential for cancer associated with exposure of law enforcement personnel to the microwave emissions from traffic radar units. A number of reports have been published in trade journals, and these have included anecdotal accounts of a small number cases of cancer, primarily ocular, skin, and testicular (Poynter 1990a, 1990b; Gibbons 1991; Clark 1991; Zaret 1991). This has been the basis for one lawsuit (*Bendure v. Kustom Signals, Inc.*), where the jury found for the radar gun manufacturer because "the officer's cancer was a

slow-developing illness that could not possibly have manifested itself only a few years after exposure, as alleged by the plaintiff" (BNA 1993). As noted in Section 3.5.6.1, Davis and Mostofi (1993) have reported an excess of observed versus expected testicular cancers in a cohort of policemen.

That these cancers could be caused by police radar units seems rather unlikely for a number of reasons. First, police radar units are low-power systems, having operational powers similar to a child's walkie-talkie (IEEE 1992b). Second, the exposure levels, as addressed in detail in Chapter 5, are very low. For example, according to Fisher (1993), the operator will be exposed at levels less than 1% of the power density at the aperture ($< 50 \mu\text{W}/\text{cm}^2$) unless: (1) there is a considerable reflection of the microwave energy, which should only occur when the reflector is closer than 365 cm (12 ft), (2) an energized unit is placed in one's lap, or (3) the operator inadvertently exposes himself. Third, the maximum anticipated exposures do not represent overexposures when compared with values recommended in state-of-the-art exposure documents. For example, if one should receive an exposure such as that described in (2) and (3), the maximum power density Fisher (1993) determined ($6.4 \text{ mW}/\text{cm}^2$) is less than the allowable human exposure criteria ($10 \text{ mW}/\text{cm}^2$) for the applicable averaging time (IEEE 1992a). Furthermore, as Fisher (1993) has pointed out, use of these devices at such close distances to the body will modify the impedance, which will reduce the radiated energy because of an impedance mismatch between the radiator and the local environment. Fourth, the penetration depth at these frequencies (10 to 35 GHz) is minimal, resulting in energy that is absorbed largely topically. Fifth, since microwaves have not been shown to be mutagenic, they should not be viewed as tumor initiators. Some animal studies have pointed to the possibility that MWs may be tumor promoters, but this needs more study. Sixth, epidemiologic studies of radar and microwave workers have not identified any statistically significant differences with cancers in general or the specific cancers mentioned ear-

lier. The radar and microwave workers included in these studies had the opportunity to be exposed at higher values of power density at a larger range of frequencies than users of traffic radar units. And last, the few animal studies that have been carried out near these frequencies have demonstrated thermal effects (Hagan and Carpenter 1976; Rosenthal et al. 1976), but at substantially higher power densities than reported for radar guns.

A similar concern has been voiced concerning brain cancer and exposure to microwave energy from low-power, portable cellular phones. This occurred after the initiation of legal activities following a single case of brain cancer in a cellular phone user. The spouse of the user brought the lawsuit based upon his perception that exposure to microwaves had caused his wife's cancer. This received national attention when the issue was discussed on the "Larry King Live" talk show on Cable News Network (Fischetti 1993).

Portable cellular phones generate about 600 mW at frequencies between 800 and 900 MHz. Although locally high values of power density may exist near the antenna, most studies have indicated that the penetration depth is low, and local SARs usually do not exceed the exposure limits (FCC 1993). However, recent studies have demonstrated differences in dosimetry, and these differences appear to depend somewhat on methodology, i.e., the use of mathematical models versus the use of tissue-equivalent phantoms, and distance from the antenna. The exact reasons for these reported differences need to be determined.

With the present state of understanding, it does not seem likely that brain cancer would be caused by exposure to these devices (FDA 1993). However, there are some significant gaps in the scientific data base, but these should be filled when well-designed animal studies and epidemiology studies are completed. At the operational frequencies, it has been demonstrated that microwave radiation produces thermal effects. Although the specific frequency band and modulation types used by portable cellular phones and other cellular devices have not been included in

studies of microwave-induced cancer in test animals, studies at 2450 MHz have not established microwaves as tumor promoters.

3.5.8.7 Summary

The incident reports indicate that overexposure to high levels of radio-frequency radiation may result in nerve damage and skin damage. It is possible that the finding of Hirsch and Parker (1952) is a case of microwave-induced cataracts, but the other reports are not consistent and do not establish a reliable link with microwave exposure.

3.5.9 Cutaneous Perception of Microwaves and RF Burns

Perception of microwave energy involves cutaneous thermal sensation or pain and demonstrates a frequency dependency at the few frequencies studied. Power densities necessary to produce threshold warmth sensations were lower at 10 GHz than at 3 GHz, for a forehead surface area of 37 cm^2 (Hendler 1968; Hendler, Hardy, and Murgatroyd 1963). This is probably due to the shallower penetration depth at the higher frequency, where more of the energy is absorbed in a smaller volume near the surface of the skin. Michaelson (1972) estimates that at 10 GHz, a thermal sensation is evoked when the entire facial skin is exposed at slightly more than 4 to $6 \text{ mW}/\text{cm}^2$ for 5 seconds or approximately $10 \text{ mW}/\text{cm}^2$ for 0.5 seconds (Michaelson 1972).

Justesen and colleagues (1982) exposed the forearm of male and female volunteers to 2.45-GHz microwaves. The SAR, estimated by measurement of the temperature in the middle of a saline-filled latex balloon, was $0.076 \text{ W}/\text{kg}/\text{mW}/\text{cm}^2$. Average thresholds of perception for women and men were $25.27 \text{ mW}/\text{cm}^2$ and $28.88 \text{ mW}/\text{cm}^2$, respectively. The values for the six participants ranged from 15.40 to $44.25 \text{ mW}/\text{cm}^2$. The researchers also exposed four of the volunteers to infrared radiation, finding that they could not tell the difference between a thermal stim-

ulus provided by the microwaves and IR radiation. A comparison between the two spectral regions showed that the perception thresholds differed by about a factor of 5, with 10.16 J of microwave energy and 1.8 J of IR radiation necessary for perception. Hence, more MW energy is necessary to perceive warmth than infrared energy.

Cook (1952) demonstrated that the threshold data for pain at 3 GHz varies from 0.83 to 3.1 W/cm², dependent upon the exposure duration, for an exposed area of 9.5 cm² (Cook 1952). However, according to Michaelson (1972), the perception of pain is "independent of the area of exposure, radiation intensity, exposure time, and anatomical site" and is related to a critical skin temperature.

Gandhi and Chatterjee (1982) determined the magnitude of the electric field that would produce short-circuit currents in common objects. Generally, the necessary E-field strength varied as a function of frequency and the object being irradiated. For example, perception (let-go in parentheses) currents were produced if a roof, fence, car, and truck were irradiated at, respectively, 250 (1040), 160 (850), 80 (440), and 20 V/m (110 V/m), between 10 to 100 kHz. The E-field strength then decreased with increasing frequency up to 10 MHz. Further work by Chatterjee, Wu, and Gandhi (1986) examined perception and let-go currents for finger and grasping contact, as discussed in Chapter 2. Average threshold perception currents for men were around 40 mA (~50 mA for pain) for finger contact from 0.1 to 3 MHz. Perception thresholds for women were lower than those for men. For example, with grasping contact, women were around 95 mA from 0.1 to 3 MHz, while men were around 120 mA. Ten-year-old children were estimated at about 60 percent of the values for male adults.

In a study of shipboard RF, Rogers (1981) found an average RF burn hazard threshold current around 200 mA between 2 and 20 MHz. According to Osepchuk (1992) "For body impedance 500 Ω , the burn threshold corresponds to 20 watts dissipated in a few seconds in a volume of 1 cc."

Comparison with "conventional" burns, Budd (1985, 1992) notes that RF burns may involve deeper body tissues because of the greater penetration depth. Clinicians have observed that undamaged layers of fat may be sandwiched between damaged layers of skin and muscle. Hot spots may be formed, producing areas of high local absorption leading to local tissue necrosis. It is possible that there may be a latency period between exposure and development of effects. Although the picture is not clear, Budd notes that there may be some clinical differences between RF burns and conventional burns.

3.6 OTHER POTENTIAL HAZARDS

3.6.1 Metallic Implants or Objects

A few studies have evaluated the interaction of radio-frequency and microwave energies with metal implants or metal objects worn by people. A list of some of these objects is in Table 3-16. Metallic objects may perturb the field, concentrating the RF energy in the vicinity of the conductive object. This may lead to localized heating, such as that observed by Feucht, Richardson, and Hines (1949). These researchers found that wire implants and metal plates increased measurable temperatures in *in vitro* and *in vivo* experiments with

Table 3-16. Devices That May Enhance or Be Affected by RF Fields

Cardiac valves
Pacemakers
Metal staples
Insulin infusion systems
Orthopedic wires, plates, and rods
Transdermal delivery system
Cochlear implants
Metal-framed spectacles
Jewelry, watches

2.45-GHz microwaves. Larger temperature increases were observed when metals were closer to the surface and with larger metal plates. Murray (1984a) reported an incident where a man wearing a transdermal nitro patch received a second degree burn the size and configuration of the patch. Apparently this occurred when energy from a recently serviced microwave oven heated the patch's aluminumized plastic adhesive strip. A carpenter with a steel plate in an index finger used "a diathermy machine which was used to dry glue on lumber." This resulted in pain and swelling when the finger was exposed (Merckel 1972).

Hocking, Joyner, and Fleming (1991) examined the interaction of RF fields with metal implants, including both heating and interference. Interference has been observed in cardiac pacemakers and cochlear implants. A number of reports and reviews are available, although with the evolution in design of pacemakers and RF sources, the reports are primarily of historical interest (Elmqvist 1976; Reis 1979). Sources of interference that have been identified include microwave ovens (Bonney, Rustan, and Ford 1973; King et al. 1970; Rustan, Hurt, and Mitchell 1973), microwaves in general (Osepchuk 1971), radar (Bonney, Rustan, and Ford 1973; Rohl et al. 1975), and diathermy units (Marshall, Scott and Peaston 1990). Obviously, interference is of great concern in the health care environment where there are numerous devices that produce RF and ELF fields, and there is a relatively high density of individuals at risk.

In the RF part of the spectrum, pacemaker interference is usually controlled by signal filtering and shielding the device. However, interference is still possible. With the complexity of today's pacemakers and RF sources, a thorough evaluation usually requires a case-by-case approach. Factors that must be considered include the operational frequency of the RF device, whether or not the fields are pulsed, and the design of the pacemaker (Grant 1993a). This approach is amenable to the industrial environment where there are usually few individuals at potential risk, but it may not be a viable approach in a

hospital. To illustrate the importance of proper selection of the pacemaker and electrode, Grant (1993b) has discussed two scenarios where pacemaker wearers experienced relatively high exposures to low-frequency magnetic fields and RF fields, with no interference.

Hepfner and Skelly (1985) reported that patients with cochlear implants "have experienced some false auditory and temperature sensations when in the presence of electromagnetic fields." RF devices and processes include two-way radios, police radar, computer terminals, and lightning. The authors also observed that interference may occur from other electrical devices that generate sub-RF frequencies, such as light dimmers, refrigerator motors, and "electrical noise generated in a therapist's mouth by metal electrolysis in her recent dental work."

3.6.2 Metal-Rimmed Spectacles

Davias and Griffin (1989) found that metal-framed spectacles perturb microwave fields between 2 and 12 GHz. They determined this by locating a field probe (monopole antenna on a ground plane) near a human phantom composed of lossy materials that simulate tissues. The reception pattern of the probe was compared when the phantom was fitted with eyewear and when the phantom wore no eyewear. They demonstrated a difference in the radiation patterns at the field probe, where there was an increase in signal intensity when the spectacles were mounted on the phantom. Their analysis indicated that the resonant frequency of the eyewear was 900 MHz and that each of the spectacle components (wing, hook, and rim) is resonant between 1.4 and 3.75 GHz.

3.6.3 Explosion Hazards

The interaction of RF with electro-explosive devices (EED) and flammable materials may

result in the ignition of materials or explosions. Materials such as solvents, fuels, and flammable gases may be ignited if the RF intensity is sufficiently high, there is a receiver of the RF energy (some conductive structure), and there is the potential for spark discharge of sufficient energy into a flammable atmosphere. In structures such as piping, stacks, and cranes, a spark discharge may occur if there is "a discontinuity in the structure which provides an intermittent electrical contact" (Maddocks 1992). For EED activation, the RF field must be sufficiently intense, and the EED must be in a position to absorb energy from the RF field (Sastradipradja and Joyner 1990). Obviously, secondary to the event, RF-induced explosions or ignitions may produce physical trauma and property damage.

Osephchuk (1992) has pointed out that a hazard unique to microwave heating is superheating of small objects. This may occur when a material is heated above the boiling point, with a rapid expansion of the superheated material resulting in a small explosion.

3.7 CONCLUSIONS

This review of the biologic-effects literature indicates there is a good deal of information supporting the existence of thermal effects of radio-frequency and microwave radiation, including cataract formation, changes in animal behavior, and effects on reproduction and development. However, even well-established thermal effects may appear to be equivocal when experiments are performed near the threshold of effects. Although the data base has grown substantially over the past decades, much of the information concerning nonthermal effects is generally inconclusive, incomplete, and sometimes contradictory. Studies of human populations have not demonstrated any reliably affected end point.

Some general limitations of research on the biologic effects of RF fields are listed in Table 3-17. Additionally, the Advisory Committee on Nonthermal Effects of Nonionizing

Table 3-17. Limitations in Biologic Effects Studies

- Differences in experimental methods
- Lack of replication of results
- Lack of establishment of mechanisms for measured end points
- Predominantly far-field experiments; relatively few data in the near field
- Animal data: primarily 915 & 2450 MHz
- Rodents used to model thermal stress are fur-bearing and do not sweat
- Epidemiologic data: no measured dose or exposure level
- Different approaches—Eastern scientists: field-based & clinical evaluations; Western scientists: primarily animal studies

Radiation of the National Research Council has offered constructive criticism of research in this field (National Research Council 1986).

3.8 REFERENCES

- Adey, W.R. 1980. Frequency and power windowing in tissue interactions with weak electromagnetic fields. *Proc. IEEE* 68: 119-125.
- Adey, W.R. 1988. Electromagnetic fields, cell membrane amplification and cancer promotion. In *Nonionizing Electromagnetic Radiations and Ultrasound* (NCRP Proceedings No. 8), pp. 88-110. Bethesda, MD: National Council on Radiation Protection and Measurements.
- Adey, W.R. and S.M. Bawin. 1982. Binding and release of brain calcium by low-level electromagnetic fields: a review. *Radio Sci.* 17(5S): 149S-157S.
- Adey, W., S. Bawin, and A. Lawrence. 1982. Effects of weak amplitude-modulated microwave fields on calcium efflux from awake cat cerebral cortex. *Bioelectromagnetics* 3: 295-307.
- Ahlborg, A. et al. 1990. Interpretation of "negative" studies in occupational epidemiology. *Scand. J. Work Environ. Health* 16: 153-157.
- Ahlborg, G. et al. 1989. Pregnancy outcome among working women. *Scand. J. Work Environ. Health* 15: 227-233.
- Alkyl, Y. et al. 1991. Immediate post-exposure effects of high-peak-power microwave pulses on operant behavior of wistar rats. *Bioelectromagnetics* 12: 183-195.
- Albert, E. 1977. Light and electron microscopic observations on the blood-brain barrier after microwave irradiation. In *Symposium on Biological Effects and Measurement of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D. Hazzard, ed., pp. 294-304. Rockville, MD: Bureau of Radiological Health.
- Albert, E.N. and M. DeSantis. 1976. Histological observations on central nervous system. In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds., pp. 299-310. Rockville, MD: Bureau of Radiological Health.
- Albert, E.N. and J.M. Kerns. 1981. Reversible microwave effects on the blood-brain barrier. *Brain Res.* 230: 153-164.
- Albrecht, R.M. and E. Landau. 1978. Microwave radiation: an epidemiological assessment. *Rev. Environ. Health* III: 43-57.
- AMA. 1987. Health effects of video display terminals. *JAMA* 257: 1508-1512.
- Anderson, L.E. 1990. ELF: Exposure levels, bio-effects and epidemiology. Presented at 1990 Health Physics Society Summer School: Assessing Non-Ionizing Radiation Hazards, June 17-22, 1990, at California State University, Fullerton, CA.
- ANSI. 1982. *Safety Levels with Respect to Human Exposure to Radio Frequency Electromagnetic Fields, 300 kHz to 100 GHz*. (ANSI C95.1-1982). New York: ANSI.
- Appleton, B. 1973. *Results of Clinical Surveys for Microwave Ocular Effects* (DHEW Pub. No. [FDA] 73-8031). Rockville, MD: Bureau of Radiological Health.
- Appleton, B. 1974a. Experimental microwave ocular effects. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 186-188. Warsaw, Poland: Polish Medical Publishers.
- Appleton, B. 1974b. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2037-2038.
- Appleton, B. et al. 1975. Microwave lens effects in humans. II. results of five-year survey. *Arch. Ophthalmol.* 93: 257-258.
- Appleton, B., S.E. Hirsch, and P.V.K. Brown. 1975. Investigation of single-exposure microwave ocular effects at 3000 MHz. *Ann. N.Y. Acad. Sci.* 247: 125-134.
- Appleton, B. and G.C. McCrossan. 1972. Microwave lens effects in humans. *Arch. Ophthalmol.* 88: 259-262.
- Archimbaud, E. et al. 1989. Acute myelogenous leukemia following exposure to microwaves. *Br. J. Haematol.* 73: 272-273.
- Archimbaud, E. et al. 1990. Correspondence. *Br. J. Haematol.* 76: 313.
- Ashani, Y., F. Henry, and G. Catravas. 1980. Combined effects of anticholinesterase drugs and low-level microwave radiation. *Radiat. Res.* 84: 496-503.
- Aurell, E. and B. Tengroth. 1973. Lenticular and retinal changes secondary to microwave exposure. *Acta. Ophthalmol.* 51: 764-769.
- Balcer-Kubiczek, E.K. and G.H. Harrison. 1985. Evidence for microwave carcinogenesis *in vitro*. *Carcinogenesis* 6: 859-864.
- Balcer-Kubiczek, E.K. and G.H. Harrison. 1989. Induction of neoplastic transformation in C3H/10T $\frac{1}{2}$ cells by 2.45-GHz microwaves and phorbol ester. *Radiat. Res.* 117: 531-537.
- Balcer-Kubiczek, E.K. and G.H. Harrison. 1991. Neoplastic transformation of C3H/10T $\frac{1}{2}$ cells following exposure to 120-Hz modulated 2.45-GHz microwaves and phorbol ester tumor promoter. *Radiat. Res.* 126: 65-72.
- Baldwin, M., S.A. Bach, and S.A. Lewis. 1960. Effects of radio-frequency energy on primate cerebral activity. *Neurology* 10: 178-187.
- Baranski, S. 1971. Effect of chronic microwave irradiation on the blood forming system of guinea pigs and rabbits. *Aerospace Med.* 42: 1196-1199.
- Baranski, S. 1972. Histological and histochemical effect of microwave irradiation on the central nervous system of rabbits and guinea pigs. *Am. J. Phys. Med.* 51: 182-191.
- Baranski, S. and P. Czerski. 1976. *Biological Effects of Microwaves*. Stroudsburg, PA: Dowden, Hutchinson & Ross, Inc.
- Baranski, S. and Z. Edelwejn. 1968. Studies on the combined effect of microwaves and some drugs on bioelectric activity of the rabbit central nervous system. *Acta Physiologica Polonica* 19: 31-41.
- Baranski, S. and Z. Edelwejn. 1975. Experimental morphologic and electroencephalographic

- studies of microwave effects on the nervous system. *Ann. N.Y. Acad. Sci.* 247: 109-116.
- Baranski, S., S. Szmigielski, and J. Moneta. 1974. Effects of microwave irradiation in vitro on cell membrane permeability. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 173-177. Warsaw, Poland: Polish Medical Publishers.
- Barnes, F.S. and C.-L. Hu. 1977. Model for some nonthermal effects of radio and microwave fields on biological membranes. *IEEE Trans. Microwave Theory Tech.* MTT-25: 742-746.
- Barron, C.I. and A.A. Baraff. 1958. Medical considerations of exposure to microwaves (radar). *JAMA* 168: 1194-1199.
- Barron, C.I., A.A. Love, and A.A. Baraff. 1955. Physical evaluation of personnel exposed to microwave emanations. *Aviat. Med.* 26: 442-452.
- Bassen, H. et al. 1987. Dosimetry for high peak-power microwave ocular effects research. In *Proceedings of the Ninth Annual Conference of the IEEE Engineering in Medicine and Biology Society* (87CH2513-0), pp. 0693-0694. New York: Institute of Electrical and Electronic Engineers.
- Baum, S.J. et al. 1976. Biological measurements in rodents exposed continuously throughout their adult life to pulsed electromagnetic radiation. *Health Phys.* 30: 161-166.
- Baumann, S. et al. 1989. Lack of effects from 2000-Hz magnetic fields on mammary adenocarcinoma and reproductive hormones in rats. *Bioelectromagnetics* 10: 329-333.
- Bawin, S. and R. Adey. 1977. Calcium binding in cerebral tissue. In *Symposium on Biological Effects and Measurement of Radio Frequency / Microwaves* (HEW Publication [FDA] 77-8026), D. Hazzard, ed., pp. 305-313. Rockville, MD: Bureau of Radiological Health.
- Bawin, S.M., R.J. Gavalas-Medici, and W.R. Adey. 1973. Effects of modulated very high frequency fields on specific brain rhythms in cats. *Brain Res.* 58: 365-384.
- Bawin, S.M., L.K. Kaczmarek, and W.R. Adey. 1975. Effects of modulated VHF fields on the central nervous system. *Ann. N.Y. Acad. Sci.* 247: 74-81.
- Beers, G.J. 1989. Biological effects of weak electromagnetic fields from 0 Hz to 200 MHz: a survey of the literature with special emphasis on possible magnetic resonance effects. *Mag. Res. Imag.* 7: 309-331.
- Bergqvist, U. 1984. Video display terminals and health. *Scan. J. Work Environ. Health* 10(suppl. 2).
- Bergqvist, U. and J.E. Wahlberg. 1994. Skin symptoms and disease during work with visual display terminals. *Contact Dermatitis* 30: 197-204.
- Berman, E., H. Carter, and D. House. 1980. Tests of mutagenesis and reproduction in male rats exposed to 2,450-MHz (CW) microwaves. *Bioelectromagnetics* 1: 65-76.
- Berman, E., H. Carter, and D. House. 1981. Observations of rat fetuses after irradiation with 2450-MHz (CW) microwaves. *J. Microwave Power* 16: 9-13.
- Berman, E., H. Carter, and D. House. 1982a. Observations of syrian hamster fetuses after exposure to 2450-MHz microwaves. *J. Microwave Power* 17: 107-112.
- Berman, E., H. Carter, and D. House. 1982b. Reduced weight in mice offspring after in utero exposure to 2450-MHz (CW) microwaves. *Bioelectromagnetics* 3: 285-291.
- Berman, E., H. Carter, and D. House. 1984. Growth and development of mice offspring after irradiation in utero with 2,450-MHz microwaves. *Teratology* 30: 393-402.
- Berman, E. et al. 1992. Fetal and maternal effects of continual exposure of rats to 970-MHz circularly-polarized microwaves. *Electro-Magnetobiol.* 11: 43-54.
- Berman, E. et al. 1985. Lethality in mice and rats exposed to 2450 MHz circularly polarized microwaves as a function of exposure duration and environmental factors. *J. Appl. Toxicol.* 5: 23-31.
- Berman, E., J. Kinn, and H. Carter. 1978. Observations of mouse fetuses after irradiation with 2.45 GHz microwaves. *Health Phys.* 35: 791-801.
- Birenbaum, L. et al. 1969a. Effect of microwaves on the rabbit eye. *J. Microw. Power* 4: 232-243.
- Birenbaum, L. et al. 1969b. Effect of microwaves on the eye. *IEEE Trans. Bio-Med. Eng. BME-16*: 7-14.
- Blackman, C.F. et al. 1985. A role for the magnetic field in the radiation-induced efflux of calcium ions from brain tissue in vitro. *Bioelectromagnetics* 6: 327-337.
- Blackman, C.F. et al. 1989. Multiple power-density windows and their possible origin. *Bioelectromagnetics* 10: 115-128.

- Blackman, C.F. 1990. ELF effects on calcium homeostasis. In *Extremely Low Frequency Electromagnetic Fields: The Question of Cancer*, B.W. Wilson, R.G. Stevens, and L.E. Anderson, eds., pp. 187-208. Columbus, OH: Battelle Press.
- Blackman, C.F. et al. 1990. Importance of alignment between local DC magnetic field and an oscillating magnetic field in responses of brain tissue in vitro and in vivo. *Bioelectromagnetics* 11: 159-167.
- Blackman, C.F., S.G. Benane, and D.E. House. 1991. The influence of temperature during electric- and magnetic-field-induced alteration of calcium-ion release from in vitro brain tissue. *Bioelectromagnetics* 12: 173-182.
- Blackman, C.F. et al. 1979. Induction of calcium-ion efflux from brain tissue by radio-frequency radiation: effects of modulation frequency and field strength. *Radio Sci.* 14(6S): 93-98.
- Blackwell, R.P. and R.D. Saunders. 1986. The effects of low-level radiofrequency and microwave radiation on brain tissue and animal behaviour. *Int. J. Radiat. Biol.* 50: 761-787.
- Blackwell, R. and A. Chang. 1988. Video display terminals and pregnancy: a review. *Br. J. Obstet. Gynecol.* 95: 446-453.
- BNA. 1993. Defense verdict issued in radar gun case: jury found no link with policeman's cancer. *Occupational Safety & Health Reporter*, p. 1554. Washington, DC: Bureau of National Affairs.
- Bollinger, J.N., R.L. Lawson, and W.C. Dolle. 1974. *Research on Biological Effects of VLF Band Electromagnetic Radiation* (Report No. SAM-TR-74-52). Brooks Air Force Base, Texas: USAF School of Aerospace Medicine. Cited in *Critique of the Literature on Bioeffects of Radiofrequency Radiation: A Comprehensive Review Pertinent to Air Force Operations* (Report No. USAFSAM-TR-87-3) by L. Heynick, p. 197. Brooks Air Force Base, TX: USAF School of Aerospace Medicine, 1987.
- Bonasera, S., J. Toler, and V. Popovic. 1988. Long-term study of 435 MHz radio-frequency radiation on blood-borne end points in cannulated rats. *J. Microw. Power Electromagn. Energy* 23: 95-104.
- Bonney, C.H., P.L. Rustan and G.E. Ford. 1973. Evaluation of effects of the microwave oven (915 and 2450 MHz) and radar (2810 and 3050 MHz) electromagnetic radiation on noncompetitive cardiac pacemakers. *IEEE Trans. Biomed. Eng. BME-20*: 357-364.
- Bonomi, L. and R. Bellucci. 1989. Considerations of the ocular pathology in 30,000 personnel of the Italian telephone company (SIP) using VDTs. *Bollettino Di Oculistica* 68 (S7): 85-98.
- Boos, S.R. et al. 1985. Work with video display terminals among office employees III. ophthalmologic factors. *Scand. J. Work Environ. Health* 11: 475-481.
- Bouchat, J.A. 1974. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2038-2039.
- Braithwaite, L. et al. 1991. Exposure of fertile chicken eggs to microwave radiation (2.45 GHz, CW) during incubation: technique and evaluation. *J. Microw. Power Electromagn. Energy* 26: 206-214.
- Brandt, L. and C.V. Nielsen. 1990. Congenital malformations among children of women working with video display terminals. *Scand. J. Work Environ. Health* 16: 329-333.
- Bridges, J.E., and M. Preache. 1981. Biological influences of power frequency electric fields—a tutorial review from a physical and experimental viewpoint. *Proc. IEEE* 69: 1092-1120.
- Brodtkin, R.H. and J. Bleiberg. 1973. Cutaneous microwave injury. *Acta Dermatol.* 53: 50-52.
- Brown-Woodman, P.D.C. et al. 1986. Radiofrequency (RF) radiation as a teratogen. *Teratology* 33: 5B.
- Brown-Woodman, P.D.C. et al. 1989. Evaluation of reproductive function of female rats exposed to radiofrequency fields (27.12 MHz) near a shortwave diathermy device. *Health Phys.* 56: 521-525.
- Brown-Woodman, P.D.C. and J.A. Hadley. 1988. Studies of the teratogenic potential of exposure of rats to 27.12 MHz pulsed short-wave radiation. *J. Bioelectricity* 7: 57-67.
- Bryant, H. and E.J. Love. 1989. Video display terminal use and spontaneous abortion risk. *Int. J. Epidemiol.* 18: 132-138.
- Budd, R.A. 1985. Can microwave/radio-frequency radiation (RFR) burns be distinguished from conventional burns? *J. Microw. Power* 20: 9-11.
- Budd, R.A. 1992. Burns associated with the use of microwave ovens. *J. Microw. Power Electromagn. Energy* 27: 160-163.
- Budd, R. and P. Czerski. 1985. Modulation of mammalian immunity by electromagnetic radiation. *J. Microw. Power* 20: 217-231.

- Butler, D.A. 1986. Do spontaneous abortion clusters among VDT operators signal a problem? In *Proceedings of the International Scientific Conference: Work with Display Units*, pp. 657-660. Stockholm, May 12-15, 1986.
- Byus, C.V. et al. 1984. Alterations in protein kinase activity following exposure of cultured human lymphocytes to modulated microwave fields. *Bioelectromagnetics* 5: 341-351.
- Byus, C.V. et al. 1988. Increased ornithine decarboxylase activity in cultured cells exposed to low energy modulated microwave fields and phorbol ester tumor promoters. *Cancer Res.* 48: 4222-4226.
- Carpenter, R.L. 1974. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2035-2036.
- Carpenter, R.L. 1975. Ocular effects of microwave radiation. *Bull. N.Y. Acad. Med.* 55: 1048-1057.
- Carpenter, R. and C. Van Ummersen. 1968. The action of microwave radiation on the eye. *J. Microw. Power* 3: 3-19.
- Carpenter, R.L. and E.M. Livstone. 1971. Evidence for nonthermal effects of microwave radiation: abnormal development of irradiated insect pupae. *IEEE Trans. Microw. Theory Tech.* MTT-19: 173-178.
- Case, R.M. 1980. "Cell calcium"—a new international, interdisciplinary journal. *Cell Calcium* 1: 1-5.
- Castren, J. et al. 1982. On the microwave exposure. *Acta Ophthalmol.* 60: 647-654.
- CDRH. 1985. *Behavioral Effects of Microwave Radiation Absorption* (HHS Publication FDA 85-8238), J.C. Monahan and J.A. D'Andrea, eds. Washington, DC: U.S. Government Printing Office.
- Chatterjee, A., R.C. Milton, and S. Thyle. 1982. Prevalence and aetiology of cataract in Punjab. *Br. J. Ophthalmol.* 66: 35-42.
- Chatterjee, I., D. Wu, and O.P. Gandhi. 1986. Human body impedance and threshold currents for perception and pain for contact hazard analysis in the VLF-MF band. *IEEE Trans. Biomed. Eng.* BME-33: 486-494.
- Chen, T.T. et al. 1988. Cataract and health status: a case-control study. *Ophthalmic. Res.* 20: 1-9.
- Chernoff, N. et al. 1987. The potential relationship of maternal toxicity, general stress, and fetal outcome. *Teratogenesis Carcinog. Mutagen* 7: 241-253.
- Chernoff, N., J.M. Rogers, and R.J. Kavlock. 1989. An overview of maternal toxicity and prenatal development: considerations for developmental toxicity hazard assessments. *Toxicology* 59: 111-125.
- Chernovetz, M.E. et al. 1975. Teratology, survival, and reversal learning after fetal irradiation of mice by 2450-MHz microwave energy. *J. Microw. Power* 10: 391-409.
- Chernovetz, M.E., D.R. Justesen, and A.F. Oke. 1977. A teratological study of the rat: microwave and infrared radiations compared. *Radio Sci.* 12(6S): 191-197.
- Chiang, H. 1988. Microwave male contraception studies in China. In *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society*. (IEEE 88CH2566-8), p. 1883. New York: Institute of Electrical and Electronic Engineers.
- Chiang, H. and B. Shao. 1989. Biological responses to microwave radiation: reproduction, development and immunology. In *Electromagnetic Interaction with Biological Systems*, J.C. Lin, ed., pp. 141-163. New York: Plenum Press.
- Chiang, H. et al. 1989. Health effects of environmental electromagnetic fields. *J. Bioelectricity* 8: 127-131.
- Chizhenkova, R.A. 1988. Slow potentials and spike unit activity of the cerebral cortex of rabbits exposed to microwaves. *Bioelectromagnetics* 9: 337-345.
- Chou, C.-K. et al. 1985. *Effects of Long-Term Radiofrequency Radiation on Immunological Competence and Metabolism* (Report No. USAF-SAM-TR-85-105). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Chou, C.-K. et al. 1992. Long-term, low-level microwave irradiation of rats. *Bioelectromagnetics* 13: 469-496.
- Chou, C.-K., A.W. Guy, and R. Galambos. 1982. Auditory perception of radio-frequency electromagnetic fields. *J. Acoust. Soc. Am.* 71: 1321-1334.
- Chou, C.-K., A.W. Guy and R.B. Johnson. 1983. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 3. SAR in Rats Exposed in 2450-MHz Circularly Polarized Waveguide* (Report No. USAFSAM-TR-83-19). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Chou, C.-K., K.C. Yee, and A.W. Guy. 1985. Auditory response in rats exposed to 2,450 MHz electromagnetic fields in a circularly polarized waveguide. *Bioelectromagnetics* 6: 323-326.
- Ciano, M. et al. 1981. High-frequency electromagnetic radiation injury to the upper extremity: local and systemic effects. *Ann. Plast. Surg.* 7: 128-135.
- Clark, J.R. 1991. Can traffic radar lead to cancer? *Law Enforcement News* 17(332): 1, 8-9.
- Cleary, S. 1977. Biological effects of microwave and radiofrequency radiation. *Crit. Rev. Environ. Controls* 27: 121-166.
- Cleary, S. 1980. Microwave cataractogenesis. *Proc. IEEE* 68: 49-55.
- Cleary, S.F. 1989. Biological effects of radiofrequency radiation: an overview. In *Electromagnetic Biointeraction Mechanisms, Safety Standards, Protection Guides*, G. Franceschetti, O.P. Gandhi, and M. Grandolfo, eds., pp. 59-79. New York: Plenum Press.
- Cleary, S.F., L.-M. Liu, and F. Garber. 1985a. Erythrocyte hemolysis by radiofrequency fields. *Bioelectromagnetics* 6: 313-322.
- Cleary, S.F., L.-M. Liu, and F. Garber. 1985b. Viability and phagocytosis of neutrophils exposed in vitro to 100-MHz radiofrequency radiation. *Bioelectromagnetics* 6: 53-60.
- Cleary, S.F., L.-M. Liu, and R.E. Merchant. 1990. Glioma proliferation modulated in vitro by isothermal radiofrequency radiation exposure. *Radiat. Res.* 121: 38-45.
- Cleary, S.F. and B.S. Pasternack. 1966. Lenticular changes in microwave workers. *Arch. Environ. Health* 12: 23-29.
- Cleary, S.F., B.S. Pasternack, and G.W. Beebe. 1965. Cataract incidence in radar workers. *Arch. Environ. Health* 11: 179-182.
- Cogan, D.G. et al. 1958. Cataracts and ultrahigh-frequency radiation. *A.M.A. Arch. Ind. Health* 18: 299-302.
- Cohen, B.H. et al. 1977. Parental factors in Down's Syndrome: results of the second Baltimore case-control study. In *Population Cytogenetics—Studies in Humans*, E.B. Hook and I.H. Porter, eds., pp. 301-352. New York: Academic Press.
- Cohen, B.H. and A.M. Lilienfeld. 1970. The epidemiological study of mongolism in Baltimore. *Ann. N.Y. Acad. Sci.* 171: 320-327.
- Cook, H.F. 1952. The pain threshold for microwave and infra-red radiation. *J. Physiol.* 118: 1-11.
- Cox, E.A. 1984. Radiation emissions from visual display units. In *Health Hazards of VDTs?* B.G. Pearce, ed., pp. 25-37. New York: Wiley.
- Curtis, R. and M. Nichols. 1983. Nonionizing radiation. In *Environmental and Occupational Medicine*, W.N. Rom, ed., pp. 693-705. Boston: Little, Brown and Company.
- Czerska, E.M. et al. 1992. Effects of continuous and pulsed 2450-MHz radiation on spontaneous lymphoblastoid transformation of human lymphocytes in vitro. *Bioelectromagnetics* 13: 247-259.
- Czerski, P. 1975. Microwave effects on the blood-forming system with particular reference to the lymphocyte. *Ann. N.Y. Acad. Sci.* 247: 232-241.
- Czerski, P. et al. 1974. Influence of microwave radiation on the hematopoietic system. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 67-74. Warsaw, Poland: Polish Medical Publishers.
- Czerski, P., M. Siekierzyński, and A. Gidyski. 1974. Health surveillance of personnel occupationally exposed to microwaves. I. Theoretical considerations and practical aspects. *Aerospace Med.* 45: 1137-1142.
- Daily, L.E. 1943. A clinical study of the results of exposure of laboratory personnel to radar and high frequency radio. *U.S. Naval Med. Bull.* 41: 1052-1056.
- Daily, L. et al. 1948. Effects of microwave diathermy on the eye. *Am. J. Physiol.* 155: 432.
- D'Andrea, J.A. 1991. Microwave radiation absorption: behavioral effects. *Health Phys.* 61: 29-40.
- D'Andrea, J.A., B.L. Cobb, and J.O. de Lorge. 1989. Lack of behavioral effects in the rhesus monkey: high peak microwave pulses at 1.3 GHz. *Bioelectromagnetics* 10: 65-76.
- D'Andrea, J.A. et al. 1979. Physiological and behavioral effects of chronic exposure to 2450-MHz microwaves. *J. Microw. Power* 14: 351-362.
- D'Andrea, J.A. et al. 1986a. Behavioral and psychological effects of chronic 2,450-MHz microwave irradiation of the rat at 0.5 mW/cm². *Bioelectromagnetics* 7: 45-56.
- D'Andrea, J.A. et al. 1986b. Intermittent exposure of rats to 2450 MHz microwaves at 2.5 mW/cm²: behavioral and physiological effects. *Bioelectromagnetics* 7: 315-328.
- D'Andrea, J., O. Gandhi, and J.L. Lords. 1977. Behavioral and thermal effects of microwave

- radiation at resonant and nonresonant wavelengths. *Radio Sci.* 12(6S): 251-256.
- D'Andrea, J.A. and J.O. de Lorge. 1990. Behavioral effects of electromagnetic fields. In *Biological Effects and Medical Applications of Electromagnetic Energy*, O.P. Gandhi, ed., pp. 317-338. Englewood Cliffs, NJ: Prentice-Hall.
- Das, K. et al. 1991. Low dose rate modulated microwave and radiofrequency radiation can alter gene expressions: alteration of neuron specific enolase activities. *Am. J. Hum. Gen.* 49(suppl. 4): 453.
- Davias, N. and D.W. Griffin. 1989. Effect of metal-framed spectacles on microwave radiation hazards to the eyes of humans. *Med. Biol. Eng. Comput.* 27: 191-197.
- Davis, R.L. and F.K. Mostofi. 1993. Cluster of testicular cancer in police officers exposed to hand-held radar. *Am. J. Ind. Med.* 24: 231-233.
- Deichmann, W.B. et al. 1963. Effects on dogs of chronic exposure to microwave radiation. *J. Occup. Med.* 9: 418-425.
- de Lorge, J.O. 1976. The effects of microwave radiation on behavior and temperature in rhesus monkeys. In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds., pp. 158-174. Rockville, MD: Bureau of Radiological Health.
- de Lorge, J.O. 1978. Disruption of behavior in mammals of three different sizes exposed to microwaves: extrapolation to larger mammals. In *Proceedings of the 1978 Symposium on Electromagnetic Fields in Biological Systems*, pp. 215-228. Edmonton, Canada: International Microwave Power Institute.
- de Lorge, J.O. 1983. The thermal basis for disruption of operant behavior by microwaves in three animal species. In *Microwaves and Thermoregulation*, E.R. Adair, ed., pp. 379-399. New York: Academic Press.
- de Lorge, J.O. 1984. Operant behavior and colonic temperature of *Macaca mulatta* exposed to radio frequency fields at and above resonant frequencies. *Bioelectromagnetics* 5: 233-246.
- de Lorge, J.O. 1985. Effects of microwaves on schedule-controlled behavior. In *Behavioral Effects of Microwave Radiation Absorption* (HHS Publication FDA 85-8238), J.C. Monahan and J.A. D'Andrea, eds., pp. 22-35. Washington, DC: U.S. Government Printing Office.
- de Lorge, J.O. and C.S. Ezell. 1980. Observing-responses of rats exposed to 1.28- and 5.62-GHz microwaves. *Bioelectromagnetics* 1: 183-198.
- Dennis, J.A., C.R. Muirhead, and J.R. Ennis. 1991a. Epidemiological studies of exposures to electromagnetic fields: I. General health and birth outcome. *J. Radiol. Prot.* 11: 3-12.
- Dennis, J.A., C.R. Muirhead, and J.R. Ennis. 1991b. Epidemiological studies of exposures to electromagnetic fields: II. Cancer. *J. Radiol. Prot.* 11: 13-25.
- DeWitt, J. et al. 1987. Behavioral effects of chronic exposure to 0.5 mW/cm² of 2,450-MHz microwaves. *Bioelectromagnetics* 8: 149-157.
- Dickason, W.L. and J.P. Barutt. 1984. Investigation of an acute microwave-oven hand injury. *J. Hand Surg.* 9A: 132-135.
- Dickersin, K. and J.A. Berlin. 1992. Meta-analysis: state-of-the-science. *Epidemiol. Rev.* 14: 154-176.
- Dietzel, F. 1975. Effects of electromagnetic radiation on implantation and intrauterine development of the rat. *Ann. N.Y. Acad. Sci.* 247: 367-376.
- Djordjevic, Z. et al. 1979. A study of the health status of radar workers. *Aviat. Space Environ. Med.* 50: 396-398.
- Djordjevic, Z., N. Lazarevic, and V. Djokovic. 1977. Studies on the hematologic effects of long-term, low-dose microwave exposure. *Aviat. Space Environ. Med.* 48: 516-518.
- Dodge, C.H. 1969. Clinical and hygienic aspects of exposure to electromagnetic fields. In *Biological Effects and Health Implications of Microwave Radiation*, S.F. Cleary, ed., pp. 140-149. Washington, DC: U.S. Government Printing Office.
- Dodge, C.H. and Z.R. Glaser. 1977. Trends in nonionizing electromagnetic radiation bioeffects and related occupational health aspects. *J. Microw. Power* 12: 319-334.
- Doll, R. 1984. Occupational cancer: problems in interpreting human evidence. *Ann. Occup. Hyg.* 28: 291-305.
- Donaldson, D.D. 1974. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2037.
- Dougherty, J.D. et al. 1965. Evaluation of an alleged case of radiation induced cataract at a radar site. *Aerospace Med.* 36: 466-471.
- Dumanski, J.D. and M.G. Sandala. 1974. The biologic action and hygienic significance of electromagnetic fields. In *Biologic Effects and*

- Health Hazards of Microwave Radiation*, pp. 289-293. Warsaw, Poland: Polish Medical Publishers.
- Durney, C.H. et al. 1978. *Radiofrequency Radiation Dosimetry Handbook*, 2nd ed. (Report SAM-TR-78-22). Brooks Air Force Base, TX: USAF School of Aerospace Medicine.
- Durney, C.H., H. Massoudi and M.F. Iskander. 1986. *Radiofrequency Radiation Dosimetry Handbook*, 4th ed. (USAFSAM-TR-85-73). Brooks Air Force Base, TX: USAF School of Aerospace Medicine.
- Dutta, S.K. et al. 1984. Microwave radiation-induced calcium ion efflux from human neuroblastoma cells in culture. *Bioelectromagnetics* 5: 71-78.
- Dutta, S.K. et al. 1989. Radiofrequency radiation-induced calcium ion efflux enhancement from human and other neuroblastoma cells in culture. *Bioelectromagnetics* 10: 197-202.
- Earle, A.M. and C.A. Blake. 1985. Effects of chronic low-level electromagnetic radiation on estrous cyclicity and fertility in the female rat. *Anat. Rec.* 211: 360-361.
- EHD. 1983. *Investigation of Radiation Emissions from Video Display Terminals*. Ottawa, Ontario, Canada: Environmental Health Directorate.
- Elder, J. 1987. Radiofrequency radiation activities and issues: a 1986 perspective. *Health Phys.* 53: 607-611.
- Elder, J.A. and Cahill, D. ed. 1984. *Biological Effects of Radiofrequency Radiation* (Report No. EPA-600/8-83-026F), Springfield, VA: National Technical Information Service (PB85-120848).
- Elder, J.A. et al. 1989. Radiofrequency radiation. In *Nonionizing Radiation Protection* (2nd ed., WHO Regional Publications European Series No. 25), M.J. Seuss and D.A. Benwell-Morison, eds., pp. 117-173. Copenhagen: World Health Organization.
- Elmqvist, H. 1976. Pacemakers and external interference. *Acta Med. Scan.* 5596: 83-86.
- Ely, T.S. 1971. Microwave death. *JAMA* 217: 1394.
- Ely, T.S. 1985. A letter to the editor. *J. Microw. Power* 20: 137.
- Ely, T.S., D.E. Goldman, and J.Z. Hearon. 1964. Heating characteristics of laboratory animals exposed to ten-centimeter microwaves. *IEEE Trans. Bio-Med. Eng.* BME-11: 123-137.
- Emery, A.F. et al. 1975. Microwave induced temperature rises in rabbit eyes in cataract research. *J. Heat Transfer* 97: 123-128.
- Ericson, A. and B. Kallen. 1986a. An epidemiological study of work with video screens and pregnancy outcome: II. a case-control study. *Am. J. Ind. Med.* 9: 459-475.
- Ericson, A. and B. Kallen. 1986b. An epidemiological study of work with video screens and pregnancy outcome: I. a registry study. *Am. J. Ind. Med.* 9: 447-457.
- FCC. 1993. *Information on Cellular Radio and Radiofrequency Radiation*. Washington, DC: Federal Communications Commission, Office of Engineering & Technology.
- FDA. 1993. Update on cellular phones. *FDA Talk Paper* (193-7). Rockville, MD: Food and Drug Administration.
- Ferri, E.S. and G.J. Hagan. 1977. Chronic low-level exposure of rabbits to microwaves. In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds., pp. 129-142. Rockville, MD: Bureau of Radiological Health.
- Feucht, B.L., A.W. Richardson, and H.M. Hines. 1949. Effects of implanted metals on tissue hyperthermia produced by microwaves. *Arch. Phys. Med.* 30: 164-169.
- Fischetti, M. 1993. The cellular phone scare. *IEEE Spectrum* 30: 43-47.
- Fisher, P.D. 1993. Microwave exposure levels encountered by police traffic radar operators. *IEEE Trans. Electromag. Compat.* 35: 36-45.
- Fisher, P.D., Lauber, J.K., and W.A.G. Voss. 1979. The effect of low-level 2450-MHz cw microwave radiation and body temperature on early embryonic development in chickens. *Radio Sci.* 14 (6S): 159-163.
- Fleck, H. 1983. Microwave oven burn. *Bull. N.Y. Acad. Med.* 59: 313-317.
- Fleming, L.E., A.M. Ducatman, and S.L. Shalat. 1991. Disease clusters: a central and ongoing role in occupational health. *J. Occup. Med.* 33: 818-825.
- Forman, S. et al. 1982. Psychological symptoms and intermittent hypertension following acute microwave exposure. *J. Occup. Med.* 24: 932-934.
- Foster, M.R., E.S. Ferri, and G.J. Hagan. 1986. Dosimetric study of microwave cataractogenesis. *Bioelectromagnetics* 7: 129-140.
- Frankos, V.H. 1985. FDA perspectives on the use of teratology data for human risk assessment. *Fundam. Appl. Toxicol.* 5: 615-625.
- Frei, M.R. et al. 1989. Thermal and physiological responses of rats exposed to 2.45-GHz radiofrequency radiation: a comparison of E.

- and H Orientation. *Radiat. Environ. Biophys.* 28: 235-246.
- Frei, M.R. et al. 1990. Field orientation effects during 5.6-GHz radiofrequency irradiation of rats. *Aviat. Space Environ. Med.* 61: 1125-1129.
- Frei, M., J. Jauchem, and F. Heinmets. 1988. Physiological effects of 2.8 GHz radio-frequency radiation: a comparison of pulsed and continuous-wave radiation. *J. Microw. Power Electromagn. Energy* 23: 85-93.
- Frei, M.R., J.R. Jauchem, and F. Heinmets. 1989. Thermoregulatory responses of rats exposed to 9.3-GHz radiofrequency radiation. *Radiat. Environ. Biophys.* 28: 67-77.
- Frey, A. 1961. Auditory system response to radio frequency energy. *Aerospace Med.* 32: 1140-1142.
- Frey, A. 1983. Letter: Comment on "microwaves and the blood-brain barrier." *J. Bioelectricity* 2: 83-88.
- Frey, A.H. 1985. Data analysis reveals significant microwave-induced eye damage in humans. *J. Microw. Power* 20: 53-55.
- Frey, A.H. and L.S. Wesler. 1990. Interaction of psychoactive drugs with exposure to electromagnetic fields. *J. Bioelectricity* 9: 187-196.
- Frey, A., S.R. Feld, and B. Frey. 1975. Neural function and behavior: defining the relationship. *Ann. N.Y. Acad. Sci.* 247: 433-439.
- Friedman, G.D. 1974. *Primer of Epidemiology*. New York: McGraw-Hill.
- Frolen, H. and B.-M. Svedenstal. 1989. Effects of pulsed magnetic fields on foetal development in mice. In *Work With Display Units* (Second International Scientific Conference), Montreal, Canada, Annexe I (Abstract 159).
- Fucic, A. et al. 1992. X-rays, microwaves and vinyl chloride monomer: their clastogenic and aneugenic activity, using the micronucleus assay on human lymphocytes. *Mutat. Res.* 282: 265-271.
- Fukui, Y. et al. 1992. Effects of hyperthermia induced by microwave irradiation on brain development in mice. *J. Radiat. Res.* 33: 1-10.
- Galloway, W. and M. Waxler. 1977. Interaction between microwave and neuroactive compounds. In *Symposium on Biological Effects and Measurements of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D. Hazard, ed., pp. 62-67. Rockville, MD: Bureau of Radiological Health.
- Galvin, M.J. et al. 1986. Influence of pre- and postnatal exposure of rats to 2.45-GHz microwave radiation on neurobehavioral function. *Bioelectromagnetics* 7: 57-71.
- Galvin, M.J. and D.I. McRee. 1986. Cardiovascular, hematologic, and biochemical effects of acute ventral exposure of conscious rats to 2450-MHz (cw) microwave radiation. *Bioelectromagnetics* 7: 223-233.
- Gandhi, O.P., ed. 1991. *Biological Effects and Medical Applications of Electromagnetic Energy*. Englewood Cliffs, NJ: Prentice-Hall.
- Gandhi, O.P. and I. Chatterjee. 1982. Radio-frequency hazards in the VLF to MF band. *Proc. IEEE* 70: 1462-1464.
- Gandhi, O.P. et al. 1992. Specific absorption rates and induced current distributions in an anatomically based human model for plane wave exposures. *Health Phys.* 63: 281-290.
- Gao, C., D. Lu, and Q. She. 1990. The effects of VDT data entry work on operators. *Ergonomics* 33: 917-924.
- Garaj-Vrhovac, V., A. Fucic, and D. Horvat. 1992. The correlation between the frequency of micronuclei and specific chromosome aberrations in human lymphocytes exposed to microwave radiation in vitro. *Mutat. Res.* 281: 181-186.
- Gardner, E.P., S.R. Ruth, and B. Render. 1988. Job stress and the VDT clerical worker. *Hum. Sys. Management* 7: 359-365.
- Garland, F.C. et al. 1990. Incidence of leukemia in occupations with potential electromagnetic field exposure in United States Navy personnel. *Am. J. Epidemiol.* 132: 293-303.
- Garson, O.M. et al. 1991. A chromosomal study of workers with long-term exposure to radio-frequency radiation. *Med. J. Australia* 155: 289-292.
- Gellin, G.A. 1971. Effect of microwave oven on facial radiodermatitis. *JAMA* 216: 1651.
- Gibbons, M. 1991. Police probe radar, cancer risk. *Advance for Radiological Science Professionals* 4(13).
- Glaser, Z. and C. Dodge. 1982. Comments on occupational safety and health practices in the USSR and some East European countries: a possible dilemma in risk assessment of RF and microwave radiation bioeffects. In *Risk/Benefit Analysis: The Microwave Case*, N. Ste-neck, ed., pp. 53-67. San Francisco: San Francisco Press.
- Goldhaber, M.K. 1990. The risk of miscarriage and birth defects among women who use visual display terminals during pregnancy. *Reproductive Toxicology* 4: 57-60.

- Goldhaber, M.K., M.R. Polen, and R.A. Hiatt. 1988. The risk of miscarriage and birth defects among women who use visual display terminals during pregnancy. *Am. J. Ind. Med.* 13: 695-706.
- Goldoni, J. 1990. Hematological changes in peripheral blood of workers occupationally exposed to microwave radiation. *Health Phys.* 58: 205-207.
- Grant, L.J. 1993a. The use of 27 MHz equipment in hospitals. Paper presented at the IEE Colloquium on EMC and Medicine, April 28, 1993. London: The Institution of Electrical Engineers.
- Grant, L.J. 1993b. Cardiac pacemakers and electromagnetic interferences. Paper presented at the IEE Colloquium on EMC and Medicine, April 28, 1993. London: The Institution of Electrical Engineers.
- Gruenau, S.P. et al. 1982. Absence of microwave effect on blood-brain barrier permeability to [¹⁴C] sucrose in the conscious rat. *Exp. Neurol.* 75: 299-307.
- Guillet, R. and S.M. Michaelson. 1977. The effect of repeated microwave exposure on neonatal rats. *Radio Sci.* 12 (6S): 125-129.
- Guy, A.W., C.-K. Chou, and B. Neuhaus. 1983. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 2. Average SAR and SAR Distribution in Man Exposed to 450-MHz RFR* (Report No. USAFSAM-TR-83-18). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Guy, A. et al. 1975a. Microwave-induced acoustic effects in mammalian auditory systems and physical materials. *Ann. N.Y. Acad. Sci.* 247: 194-218.
- Guy, A. et al. 1975b. Effect of 2450-MHz radiation on the rabbit eye. *IEEE Trans. Microw. Theory Tech.* MTT-23: 492-498.
- Guy, A.W. et al. 1980. Long-term 2450-MHz microwave irradiation of rabbits: methodology and evaluation of ocular and physiologic effects. *J. Microw. Power* 15: 37-44.
- Guy, A.W. et al. 1983. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 1. Design, Facilities, and Procedures* (Report No. USAFSAM-TR-83-17). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Guy, A.W. et al. 1985. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 9. Summary* (Report No. USAF-SAM-TR-85-64). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Hagaman, B.G. 1989a. "Submission of revised work procedures for Haiku site" (letter to H. Tateishi). Alexandria, VA: Kershner, Wright & Hagaman, P.C.
- Hagaman, B.G. 1989b. "Electrostatic test meters for DOT Haiku contractors" (letter to H. Tateishi). Alexandria, VA: Kershner, Wright & Hagaman, P.C.
- Hagan, G.J. and R.L. Carpenter. 1976. Relative cataractogenic potencies of two microwave frequencies (2.45 and 10 GHz). In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds., pp. 143-155. Rockville, MD: Bureau of Radiological Health.
- Hamburger, S., J.N. Logue, and P.M. Silverman. 1983. Occupational exposure to non-ionizing radiation and an association with heart disease: an exploratory study. *J. Chron. Dis.* 36: 791-802.
- Hamrick, P.E. and D.I. McRee. 1975. Exposure of the Japanese quail embryo to 2.45 GHz microwave radiation during the second day of development. *J. Microw. Power* 10: 211-221.
- Hamrick, P.E. and S.S. Fox. 1977. Rat lymphocytes in cell culture exposed to 2450 MHz (CW) microwave radiation. *J. Microw. Power* 12: 125-132.
- Harding, J.J., R.S. Harding, and M. Egerton. 1989. Risk factors for cataract in Oxfordshire: diabetes, peripheral neuropathy, myopia, glaucoma, and diarrhoea. *Acta Ophthalmol.* 67: 510-517.
- Haseman, J.K. and M.D. Hogan. 1975. Selection of the experimental unit in teratology studies. *Teratology* 12: 165-172.
- Healer, J. 1969. Review of studies of people occupationally exposed to radio-frequency radiations. In *Biological Effects and Health Implications of Microwave Radiation*, S.F. Cleary, ed., pp. 90-97. Washington, DC: U.S. Government Printing Office.
- Heins, A.P. 1990. "RF investigation involving the Maersk Constellation" (letter to M. Grueber). Salt Lake City, UT: USDOL OSHA Health Response Team.
- Heins, A.P. and R. Curtis. 1990. "RF shock hazard inspection terminal 5, Port of Seattle" (letter to D. Beeston). Salt Lake City, UT: USDOL OSHA Health Response Team.
- Hendler, E. 1968. Cutaneous receptor response to microwave radiation. In *Thermal Problems*

- U.S. Environmental Protection Agency, Health Effects Research Laboratory, Research Triangle Park, NC. Cited in EPA 1984. *Biological Effects of Radiofrequency Radiation* (Report No. EPA-600/8-83-026F), J. Elder and D. Cahill, ed., Springfield, VA: National Technical Information Service (PB85-120848).
- Kaplan, J. et al. 1982. Biological and behavioral effects of prenatal and postnatal exposure to 2450-MHz electromagnetic radiation in the squirrel monkey. *Radio Sci.* 17(5S): 135S-144S.
- Kavet, R. and R. Tell. 1991. VDTs: field levels, epidemiology and laboratory studies. *Health Phys.* 61: 47-57.
- King, G.R. et al. 1970. Effect of microwave oven on implanted cardiac pacemaker. *JAMA* 212: 1213.
- Kirk, W.P. 1984. Life span and carcinogenesis. In *Biological Effects of Radiofrequency Radiation* (Report No. EPA-600/8-83-026F), J. Elder and D. Cahill, eds., pp. 5-106-5-111. Springfield, VA: National Technical Information Service (PB85-120848).
- Klaunberg, B.J. and J.H. Merritt. 1991. No evidence for microwave interaction with noradrenergic neurotransmitter systems. *Health Phys.* 60: 282-283.
- Klimkova-Deutschova, E. 1974. Neurologic findings in persons exposed to microwaves. In *Biological Effects and Health Hazards of Microwave Radiation*, pp. 268-272. Warsaw, Poland: Polish Medical Publishers.
- Kolmodin-Hedman, B. et al. 1988. Health problems among operators of plastic welding machines and exposure to radiofrequency electromagnetic fields. *Int. Arch. Occup. Environ. Health* 60: 243-247.
- Kondra, P.A. et al. 1970. Growth and reproduction of chickens subjected to microwave radiation. *Can. J. Anim. Sci.* 50: 639-644.
- Kramar, P., C. Harris, A.W. Guy. 1987. Thermal cataract formation in rabbits. *Bioelectromagnetics* 8: 397-406.
- Krause, D. et al. 1991. Microwave exposure alters the expression of 2.5-A-dependent RNase. *Radiat. Res.* 127: 164-170.
- Kues, H.A. and S. D'Anna. 1987. Changes in the monkey eye following pulsed 2.45-GHz microwave exposure. *Proceedings of the Ninth Annual Conference of the IEEE Engineering in Medicine and Biology Society* (vol. 2; IEEE Cat. No. 87CH2513-0), pp. 698-700. New York: Institute of Electrical and Electronic Engineers.
- Kues, H.A. et al. 1985. Effects of 2.45-GHz microwaves on primate corneal endothelium. *Bioelectromagnetics* 6: 177-188.
- Kues, H.A. et al. 1992. Increased sensitivity of the non-human primate eye to microwave radiation following ophthalmic drug pretreatment. *Bioelectromagnetics* 13: 379-393.
- Kues, H.A. and J.C. Monahan. 1992. Microwave-induced changes to the primate eye. *Johns Hopkins APL Tech. Dig.* 13: 244-254.
- Kunz, L.L. et al. 1983. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 5. Evaluation of the Immune System's Response* (Report No. USAFSAM-TR-83-50). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Kunz, L.L. et al. 1984. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 6. Hematological, Serum Chemistry, Thyroxine, and Protein Electrophoresis Evaluations* (Report No. USAFSAM-TR-84-2). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Kunz, L.L. et al. 1985. *Effects of Long-Term Low-Level Radiofrequency Radiation Exposure on Rats Volume 8. Evaluation of Longevity, Cause of Death, and Histopathological Findings* (Report No. USAFSAM-TR-85-11). Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine.
- Kurppa, K. et al. 1985. Birth defects and exposure to video display terminals during pregnancy. *Scand. J. Work Environ. Health* 11: 353-356.
- Kurt, T.L. 1988. Re: "Increased mortality in amateur radio operators due to lymphatic and hematopoietic malignancies." *Am. J. Epidemiol.* 128: 1384.
- Lai, H. et al. 1984a. Effects of acute low-level microwaves on pentobarbital-induced hypothermia depend on exposure orientation. *Bioelectromagnetics* 5: 203-211.
- Lai, H. et al. 1984b. Ethanol-induced hypothermia and ethanol consumption in the rat are affected by low-level microwave irradiation. *Bioelectromagnetics* 5: 213-220.
- Lai, H. et al. 1987. A review of microwave irradiation and actions of psychoactive drugs. *IEEE Eng. Med. Biol. Mag.* 6: 31-36.
- Lai, H. 1992. Research on the neurological effects of nonionizing radiation at the University of Washington. *Bioelectromagnetics* 13: 513-526.

- Lancranjan, I. et al. 1975. Gonadic function in workmen with long-term exposure to microwaves. *Health Phys.* 29: 381-383.
- LaRoche, L.P., M.M. Zaret, and A.F. Braun. 1970. An operational safety program for ophthalmic hazards of microwave. *Arch. Environ. Health* 20: 350-355.
- Larsen, A.I. 1991. Congenital malformations and exposure to high-frequency electromagnetic radiation among Danish physiotherapists. *Scand. J. Work Environ. Health* 17: 318-323.
- Larsen, A.I., J. Olsen, and O. Svane. 1991. Gender-specific reproductive outcome and exposure to high-frequency electromagnetic radiation among physiotherapists. *Scand. J. Work Environ. Health* 17: 324-329.
- Larsen, A.I. and J. Skotte. 1991. Can exposure to electromagnetic radiation in diathermy operators be estimated from interview data?: a pilot study. *Am. J. Ind. Med.* 19: 51-57.
- Lary, J.M. 1991. "Interpretation of teratological results in Tofani et al. 1986." October 1, 1991 (personal communication). J.M. Lary, Center for Disease Control, 1600 Clifton Road, Atlanta, GA 30333.
- Lary, J.M. and D.L. Conover. 1987. Teratogenic effects of radiofrequency radiation. *IEEE Eng. Med. Biol. Mag.* 6: 42-46.
- Lary, J.M., D.L. Conover, and P.H. Johnson. 1983. Absence of embryotoxic effects from low-level (nonthermal) exposure of rats to 100 MHz radiofrequency radiation. *Scand. J. Work Environ. Health* 9: 120-127.
- Lary, J.M. et al. 1982. Teratogenic effects of 27.12 MHz radiofrequency radiation in rats. *Teratology* 26: 299-309.
- Lary, J.M. et al. 1983. Teratogenicity of 27.12-MHz radiation in rats is related to duration of hyperthermic exposure. *Bioelectromagnetics* 4: 249-255.
- Lary, J.M. et al. 1986. Dose-response relationship and birth defects in radiofrequency-irradiated rats. *Bioelectromagnetics* 7: 141-149.
- Lebovitz, R.M. 1981. Prolonged microwave irradiation of rats: effects on concurrent operant behavior. *Bioelectromagnetics* 2: 169-185.
- Lebovitz, R.M. and L. Johnson. 1987. Acute, whole-body microwave exposure and testicular function of rats. *Bioelectromagnetics* 8: 37-43.
- Lester, J.R. 1985. Reply to "cancer mortality and air force bases: a reevaluation." *J. Bioelectricity* 4: 129-131.
- Lester, J.R. and D.F. Moore. 1982a. Cancer incidence and electromagnetic radiation. *J. Bioelectricity* 1: 59-76.
- Lester, J.R. and D.F. Moore. 1982b. Cancer mortality and air force bases. *J. Bioelectricity* 1: 77-82.
- Liburdy, R.P. 1979. Radiofrequency radiation alters the immune system: modulation of T- and B-lymphocyte levels and cell-mediated immunocompetence by hyperthermic radiation. *Radiat. Res.* 77: 34-46.
- Liddle, C.G. and C.F. Blackman. 1984. Endocrine, physiological and biochemical effects. In *Biological Effects of Radiofrequency Radiation* (Report No. EPA-600/8-83-026F), J. Elder and D. Cahill, eds., pp. 5-79-5-93. Springfield, VA: National Technical Information Service (PB85-120848).
- Liddle, C.G., J.P. Putnam, and O.P. Huey. 1994. Alteration of life span of mice chronically exposed to 2.45 GHz CW microwaves. *Bioelectromagnetics* 15: 177-181.
- Lilienfeld, A.M. 1976. *Foundations of Epidemiology*. New York: Oxford University Press.
- Lilienfeld, A.M. et al. 1978. *Evaluation of Health Status of Foreign Service and Other Employees from Selected Eastern European Posts* (NTIS PB288163). Springfield, VA: National Technical Information Service.
- Lim, J.I. et al. 1993. Visual abnormalities associated with high-energy microwave exposure. *Retina* 13: 230-233.
- Lin, J.C. 1979. Health aspects of radio and microwave radiation. *J. Environ. Pathol. Toxicol.* 2: 1413-1432.
- Lin, J. 1980. The microwave auditory phenomenon. *Proc. IEEE* 68: 67-73.
- Lin, J.C. 1989. Pulsed radiofrequency field effects in biological systems. In *Electromagnetic Interaction with Biological Systems*, J.C. Lin, ed., pp. 165-177. New York: Plenum Press.
- Lin, J.C., H.I. Bassen, and C.-L. Wu. 1977. Perturbation effect of animal restraining materials on microwave exposure. *IEEE Trans. Biomed. Eng. BME-24*: 80-83.
- Lin, J.C. and M.F. Lin. 1982. Microwave hyperthermia-induced blood-brain barrier alterations. *Radiat. Res.* 89: 77-87.
- Lin, J.C., J.C. Nelson, and M.E. Ekstrom. 1979. Effects of repeated exposure to 148-MHz radio waves on growth and hematology of mice. *Radio Sci.* 14(6S): 173-179.
- Lindholm, M.-L. et al. 1992. Magnetic fields of video display terminals and spontaneous abortion. *Am. J. Epidemiol.* 9: 1041-1051.

- Litovitz, T.A. et al. 1993. The role of coherence time in the effect of microwaves on ornithine decarboxylase activity. *Bioelectromagnetics* 14: 395-403.
- Liu, Y.-H. et al. 1991. Biopsies of human testes receiving multiple microwave irradiation: an histological and ultramicroscopical study. *J. Bioelectricity* 10: 213-230.
- Liu, L.-M., F.G. Nickless, and S.F. Cleary. 1979. Effects of microwave radiation on erythrocyte membranes. *Radio Sci.* 14(6S): 109-115.
- Logue, J.N. et al. 1985. Congenital anomalies and paternal occupational exposure to short-wave, microwave, infrared, and acoustic radiation. *J. Occup. Med.* 27: 451-452.
- Lotz, W.G. and S.M. Michaelson. 1978. Temperature and corticosterone relationships in microwave-exposed rats. *J. Appl. Physiol. Respir. Environ. Exercise Physiol.* 44: 438-445.
- Lotz, W.G. and R.P. Podgorski. 1982. Temperature and adrenocortical responses in rhesus monkeys exposed to microwaves. *J. Appl. Physiol. Respir. Environ. Exercise Physiol.* 53: 1565-1571.
- Lu, S.-T. et al. 1981. Microwave-induced temperature, corticosterone, and thyrotropin interrelationships. *J. Appl. Physiol. Respir. Environ. Exercise Physiol.* 50: 399-405.
- Lu, S.-T. et al. 1992. Abnormal cardiovascular responses induced by localized high power microwave exposure. *IEEE Trans. Biomed. Eng.* 39: 484-492.
- Lu, S.-T., G. Lotz, and S.M. Michaelson. 1980. Advances in microwave-induced neuroendocrine effects: the concept of stress. *Proc. IEEE* 68: 73-77.
- Lu, S.-T. and S.M. Michaelson. 1987. Comments on "effects of continuous low-level exposure to radiofrequency radiation on intrauterine development in rats." *Health Phys.* 53: 545-546.
- Lyle, D.B. et al. 1983. Suppression of T-lymphocyte cytotoxicity following exposure to sinusoidally amplitude-modulated fields. *Bioelectromagnetics* 4: 281-292.
- Mackay, C. 1987. The alleged reproductive hazards of VDUs. *Work & Stress* 1: 49-57.
- Maddocks, A.J. 1992. The ignition of flammable atmospheres by radio frequency initiated sparks. *International Electromagnetic Compatibility Symposium Proceedings*, pp. 129-139. Singapore: Nanyang Technological University.
- Maes, A. et al. 1993. In vitro cytogenic effects of 2450 MHz waves on human peripheral blood lymphocytes. *Bioelectromagnetics* 14: 495-501.
- Magin, C. 1994. "Investigation of an unlicensed uplink station." February 15, 1994 (personal communication). Charles Magin, Federal Communications Commission, Enforcement Division, P.O. Box 250, Columbia, MD 21045.
- Majewska, K. 1968. Investigations on the effect of microwaves on the eye. *Pol. Med. J.* 38: 989-994.
- Marcus, M. 1989. Letter to the editor. *Scand. J. Work Environ. Health* 15: 156-158.
- Marcus, M. 1990. Epidemiologic studies of VDT use and pregnancy outcome. *Reprod. Toxicol.* 4: 51-56.
- Marha, K. 1971. Microwave radiation safety standards in Eastern Europe. *IEEE Trans. Microw. Theory Tech.* MTT-19: 165-168.
- Marino, A.A., R.O. Becker, and B. Ullrich. 1976. The effect of continuous exposure to low frequency electric fields on three generations of mice: a pilot study. *Experientia* 32: 565-566.
- Marino, A.A. et al. 1980. Power frequency electric field induces biological changes in successive generations of mice. *Experientia* 36: 309-311.
- Marriott, I.A. and M.A. Stuchly. 1986. Health aspects of work with visual display terminals. *J. Occup. Med.* 28: 833-848.
- Marshall, I., D.H.T. Scott, and I.A. Peaston. 1990. A diathermy suppression filter for external pacemakers. *J. Med. Eng. Technol.* 14: 155-157.
- McAfee, R.D. et al. 1979. Absence of ocular pathology after repeated exposure of unanesthetized monkeys to 9.3-GHz microwaves. *J. Microw. Power* 14: 41-44.
- McDonald, A.D. et al. 1986. Spontaneous abortion and occupation. *J. Occup. Med.* 28: 1232-1238.
- McDonald, A.D. et al. 1987. Occupation and pregnancy outcome. *Br. J. Ind. Med.* 44: 521-526.
- McDonald, A.D. et al. 1988a. Prematurity and work in pregnancy. *Br. J. Ind. Med.* 45: 56-62.
- McDonald, A.D. et al. 1988b. Fetal death and work in pregnancy. *Br. J. Ind. Med.* 45: 148-157.
- McDonald, A.D. et al. 1988c. Work with visual display units and pregnancy. *Br. J. Ind. Med.* 45: 509-515.

- McLaughlin, J.T. 1957. Tissue destruction and death from microwave radiation (radar). *Calif. Med.* 86: 336-339.
- McRee, D.I. 1980. Soviet and Eastern European research on biological effects of microwave radiation. *Proc. IEEE* 68: 84-91.
- McRee, D.I. et al. 1975. Some effects of exposure of the Japanese quail embryo to 2.45-GHz microwave radiation. *Ann. N.Y. Acad. Sci.* 247: 377-390.
- McRee, D.I. et al. 1980. Long-term 2450-MHz cw microwave irradiation of rabbits: evaluation of hematological and immunological effects. *J. Microw. Power* 15: 45-52.
- Merckel, C. 1972. Microwave and man—the direct and indirect hazards, and the precautions. *Calif. Med.* 117: 20-24.
- Merriam, G.R. 1974. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2036-2037.
- Merritt, J.H., A.F. Chamness, and S.J. Allen. 1978. Studies on blood-brain barrier permeability after microwave-radiation. *Rad. Environ. Biophys.* 15: 367-377.
- Merritt, J.H., W.W. Shelton, and A.F. Chamness. 1982. Attempts to alter $^{45}\text{Ca}^{2+}$ binding to brain tissue with pulse-modulated microwave energy. *Bioelectromagnetics* 3: 475-478.
- Michaelson, S.M. 1971. Biomedical aspects of microwave exposure. *Am. Ind. Hyg. Assoc. J.* 32: 338-345.
- Michaelson, S.M. 1972. Cutaneous perception of microwaves. *J. Microw. Power* 7: 67-73.
- Michaelson, S.M. 1974. Thermal effects of single and repeated exposures to microwaves—a review. In *Biological Effects and Health Hazards of Microwave Radiation*, pp. 1-14. Warsaw, Poland: Polish Medical Publishers.
- Michaelson, S. 1980. Microwave biological effects: an overview. *Proc. IEEE* 68: 40-49.
- Michaelson, S. 1982a. Microwave and radiofrequency radiation. In *Nonionizing Radiation Protection*, 1st ed., WHO Regional Publications European Series No. 10, M.J. Seuss, ed., pp. 145-174. Copenhagen: World Health Organization.
- Michaelson, S. 1982b. Health implications of exposures to radiofrequency/microwave energies. *Br. J. Ind. Med.* 31: 105-119.
- Michaelson, S. 1983. Analysis of experimental and epidemiological data from exposure to microwave/radiofrequency (MW/RF) energies. In *Biological Effects and Dosimetry of Nonionizing Radiation Radiofrequency and Microwave Energies*, M. Grandolfo, S. Michaelson, and A. Rindi, eds., pp. 589-609. New York: Plenum Press.
- Michaelson, S.M. 1985. Subtle effects of radiofrequency energy absorption and their physiological implications. In *Interactions between Electromagnetic Fields and Cells*, A. Chiabrera, C. Nicolini, and H.P. Schwan, eds., pp. 581-602. New York: Plenum Press.
- Michaelson, S.M. 1986. Interaction of nonmodulated radio frequency fields with living matter: experimental results. In *CRC Handbook of Biological Effects of Electromagnetic Fields*, C. Polk and E. Postow, eds., pp. 339-423. Boca Raton, FL: CRC Press.
- Michaelson, S. 1991. Biological effects of radiofrequency radiation: concepts and criteria. *Health Phys.* 61: 3-14.
- Michaelson, S.M. and C.H. Dodge. 1971. Soviet views on the biological effects of microwaves—an analysis. *Health Phys.* 21: 108-111.
- Michaelson, S. et al. 1961. Tolerance of dogs to microwave exposure under various conditions. *Ind. Med. Surg.* 30(7): 298.
- Michaelson, S.M. et al. 1964. The hematologic effects of microwave exposure. *Aerospace Med.* 35: 824-829.
- Michaelson, S.M. et al. 1977. Neuroendocrine responses in the rat and dog exposed to 2450 MHz (cw) microwaves. In *Symposium on Biological Effects and Measurement of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D. Hazzard, ed., pp. 263-279. Rockville, MD: Bureau of Radiological Health.
- Michaelson, S.M., J.W. Howland, and W.B. Deichmann. 1971. Response of the dog to 24,000 and 1285 MHz microwave exposure. *Ind. Med.* 40: 18-23.
- Michaelson, S.M. and J.C. Lin. 1987. *Biological Effects and Health Implications of Radiofrequency Radiation*. New York: Plenum Press.
- Michaelson, S.M. and J. Osepchuk. 1974. Letters of comment—pro and con. *N.Y. State J. Med.* 74: 2034-2035.
- Michaelson, S.M., R.A.E. Thomson, and J.W. Howland. 1965. Comparative studies on 1285 and 2800 Mc/sec pulsed microwaves. *Aerospace Med.* 36: 1059-1064.
- Mikolajczyk, H. 1976. Microwave-induced shifts of gonadotropic activity in anterior pituitary gland of rats. In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds.,

- pp. 377-383. Rockville, MD: Bureau of Radiological Health.
- Milham, S. 1982. Mortality from leukemia in workers exposed to electrical and magnetic fields. *N. Eng. J. Med.* 307: 249.
- Milham, S. 1985a. Mortality in workers exposed to electromagnetic fields. *Environ. Health Perspect.* 62: 297-300.
- Milham, S. 1985b. Silent keys: leukemia mortality in amateur radio operators. *Lancet* 1: 812.
- Milham, S. 1988a. Increased mortality in amateur radio operators due to lymphatic and hematopoietic malignancies. *Am. J. Epidemiol.* 127: 50-54.
- Milham, S. 1988b. The author replies. *Am. J. Epidemiol.* 128: 1384-1385.
- Milham, S. 1988c. Mortality by license class in amateur radio operators. *Am. J. Epidemiol.* 128: 1175-1176.
- Milroy, W.C. and S.M. Michaelson. 1972a. Microwave cataractogenesis: a critical review of the literature. *Aerospace Med.* 43: 67-75.
- Milroy, W.C. and S.M. Michaelson. 1972b. Thyroid pathophysiology of microwave radiation. *Aerospace Med.* 43: 1126-1131.
- Mitchell, C.L. 1985. Soviet research on microwave-behavior interactions. In *Behavioral Effects of Microwave Radiation Absorption* (HHS Publication [FDA] 85-8238), J.C. Monahan and J.A. D'Andrea, eds., pp. 1-8. Washington, DC: U.S. Government Printing Office.
- Mitchell, C.L. et al. 1988. Some behavioral effects of short-term exposure of rats to 2.45 GHz microwave radiation. *Bioelectromagnetics* 9: 259-268.
- Mitchell, C.L. et al. 1989. Results of a United States and Soviet Union joint project on nervous system effects of microwave radiation. *Environ. Health Perspect.* 81: 201-209.
- Mitchell, D.S., W.G. Switzer, and E.L. Bronaugh. 1977. Hyperactivity and disruption of operant behavior in rats after multiple exposures to microwave radiation. *Radio Sci.* 12(6S): 263-271.
- Monahan, J.C. and W.W. Henton. 1977. Microwave absorption and taste aversion as a function of 915 MHz radiation. In *Symposium on Biological Effects and Measurement of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D.G. Hazard, ed., pp. 34-40. Rockville, MD: Bureau of Radiological Health.
- Monahan, J.C. and W.W. Henton. 1979. The effect of psychoactive drugs on operant behavior induced by microwave radiation. *Radio Sci.* 14(6S): 233-238.
- Monson, R.R. 1990. *Occupational Epidemiology*, 2nd ed. Boca Raton, FL: CRC Press, Inc.
- Motzkin, S.M. 1987. Millimeter wave effects of natural and model cellular and subcellular systems. In *Proceedings of the Ninth Annual Conference of the IEEE Engineering in Medicine and Biology Society* (87CH2513-0), pp. 0446-0447. New York: Institute of Electrical and Electronic Engineers.
- Muraca, G.J., E.S. Ferri, and F.L. Buchta. 1976. A study of the effects of microwave irradiation of the rat testes. In *Biological Effects of Electromagnetic Waves* (HEW Publication [FDA] 77-8010), C.C. Johnson and M.L. Shore, eds., pp. 484-494. Washington, DC: U.S. Government Printing Office.
- Murray, K.B. 1984a. Hazard of microwave ovens to transdermal delivery system. *N. Engl. J. Med.* 310: 721.
- Murray, W.E. 1984b. Video display terminals: radiation issues. *IEEE Comp. Graphics Appl.* 4: 41-44.
- Murray, W.E. et al. 1981. A radiation and industrial hygiene survey of video display terminal operations. *Hum. Factors* 23: 413-420.
- National Research Council. 1983. *Video Displays, Work, and Vision*. Washington, DC: National Academy Press, 1983.
- National Research Council. 1986. *Nonthermal Effects of Nonionizing Radiation*. Washington, DC: National Academy Press (National Research Council).
- Nawrot, P.S., D.I. McRee, and M.J. Galvin. 1985. Teratogenic, biochemical, and histological studies with mice prenatally exposed to 2.45-GHz microwave radiation. *Radiat. Res.* 102: 35-45.
- Nawrot, P.S., D.I. McRee, and R.E. Staples. 1981. Effects of 2.45 GHz cw microwave radiation on embryofetal development in mice. *Teratology* 24: 303-314.
- NCRP. 1986. *Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields* (NCRP Report No. 86). Bethesda, MD: NCRP.
- NCRP. 1988. *Nonionizing Electromagnetic Radiations and Ultrasound* (NCRP Proceedings No. 8), pp. 9-125. Bethesda, MD: National Council on Radiation Protection and Measurements.

- Neilly, J.P. and J.C. Lin. 1986. Interaction of ethanol and microwaves on the blood-brain barrier of rats. *Bioelectromagnetics* 7: 405-414.
- Nelson, B.K. et al. 1991. Marked increase in teratogenicity of the combined administration of the industrial solvent 2-methoxyethanol and radiofrequency radiation in rats. *Teratology* 43: 621-634.
- Nielsen, C.V. and L. Brandt. 1990. Spontaneous abortion among women using video display terminals. *Scand. J. Work Environ. Health* 16: 323-328.
- Nielsen, C.V. and L. Brandt. 1992. Fetal growth, preterm birth and infant mortality in relation to work with video display terminals during pregnancy. *Scand. J. Work Environ. Health* 18: 346-350.
- Nilsson, R. et al. 1989. Microwave effects on the central nervous system—a study of radar mechanics. *Health Phys.* 56: 777-779.
- National Telecommunications and Information Administration. 1981. *Microwave Radiation at the U.S. Embassy in Moscow and Its Biological Implications: An Assessment*, (NTIA-SP-81-12). Washington, DC: U.S. Department of Commerce.
- Nurminen, T. and K. Kurppa. 1988. Office employment, work with video display terminals, and course of pregnancy. *Scand. J. Work Environ. Health* 14: 293-298.
- O'Connor, M.E. 1980. Mammalian teratogenesis and radio-frequency fields. *Proc. IEEE* 68: 56-60.
- O'Connor, M.E. 1985. Effects of radiofrequency radiation during development: the prenatal period as a model system. In *Behavioral Effects of Microwave Radiation Absorption* (HHS Publication [FDA] 85-8238), J.C. Monahan and J.A. D'Andrea, eds., pp. 102-109. Washington, DC: U.S. Government Printing Office.
- O'Connor, M.E. 1988. Behavioral effects regarding exposure to nonionizing electromagnetic radiation. In *Nonionizing Electromagnetic Radiations and Ultrasound* (Proceedings No. 8), pp. 65-79. Bethesda, MD: National Council on Radiation Protection and Measurements.
- O'Connor, M.E. 1990. Teratogenesis: nonionizing electromagnetic fields. In *Biological Effects and Medical Applications of Electromagnetic Energy*, O.P. Gandhi, ed., pp. 357-372. Englewood Cliffs, NJ: Prentice-Hall.
- Odland, L.T. 1972. Observations on microwave hazards to USAF personnel. *J. Occup. Med.* 14: 544-547.
- Olsen, R.G. 1977. Insect teratogenesis in a standing-wave irradiation system. *Radio Sci.* 12(6S): 199-207.
- Olshan, A.F., K. Teschke, and M. Baird. 1991. Paternal occupation and congenital anomalies in offspring. *Am. J. Ind. Med.* 20: 447-475.
- Oscar, K. and D. Hawkins. 1977. Microwave alteration of the blood-brain barrier system of rats. *Brain Res.* 126: 281-293.
- Oscar, K. et al. 1981. Local cerebral blood flow after microwave exposure. *Brain Res.* 204: 220-225.
- Osepchuk, J.M. 1971. Comparison of potential device interference and biological exposure hazards in microwave leakage fields. In *IEEE International Electromagnetic Compatibility Symposium Record*, 1973, pp. 155-161. New York: IEEE.
- Osepchuk, J.M. 1983. *Biological effects of Electromagnetic Radiation*. New York: IEEE.
- Osepchuk, J.M. 1992. *Radiofrequency Safety Issues in Industrial Heating Systems*. Springfield, VA: National Technical Information Service (AD-P007726/3/XAB).
- Oser, B.L. 1981. The rat as a model for human toxicological evaluation. *J. Toxicol. Environ. Health* 8: 521-542.
- Ouellet-Hellstrom, R. and W.F. Stewart. 1993. Miscarriages among female physical therapists who report using radio- and microwave-frequency electromagnetic radiation. *Am. J. Epidemiol.* 138: 775-786.
- Parazzini, F. et al. 1993. Video display terminal use during pregnancy and reproductive outcome—a meta analysis. *J. Epidemiol. Comm. Health* 47: 265-268.
- Parker, L.N. 1973. Thyroid suppression and adenomullary activation by low-intensity microwave radiation. *Am. J. Physiol.* 224: 1388-1390.
- Paulsson, L.-E., Y. Hamnerius, H.-A. Hansson, and J. Sjostrand. 1979. Retinal damage experimentally induced by microwave radiation at 55 mW/cm². *Acta Ophthalmol.* 57: 183-197.
- Pay, L., E.C. Beyer, and C.F. Reichelderfer. 1972. Microwave effects on reproductive capacity and genetic transmission in *Drosophila melanogaster*. *J. Microw. Power* 7: 75-82.
- Pazderova, J., J. Pickova, and V. Bryndova. 1974. Blood proteins in personnel of television and radio transmitting stations. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 281-288. Warsaw, Poland: Polish Medical Publishers.

- Petersen, R.C. 1983. Bioeffects of microwaves: a review of current knowledge. *J. Occup. Med.* 25: 103-111.
- Petersen, R.C. 1991. "Additional information on NIOSH vdt reproductive effects study." August 19, 1991 (personal communication). R.C. Petersen, AT&T, 6000 Mountain Avenue, Murray Hill, NJ 07974-0636.
- Peterson, D.J., L.M. Pardow, and O.P. Gandhi. 1979. An investigation of the thermal and athermal effects of microwave irradiation on erythrocytes. *IEEE Trans. Biomed. Eng.* BME-26: 428-436.
- Phillips, R.D. et al. 1975. Thermoregulatory, metabolic, and cardiovascular responses of rats to microwaves. *J. Appl. Physiol.* 38: 630-635.
- Polifka, J.E. and J.M. Friedman. 1991. Environmental toxins and recurrent pregnancy loss. *Infertility and Reproductive Medicine Clinics of North America* 2: 195-213.
- Polk, C. and E. Postow, ed. 1986. *Handbook of Biological Effects of Electromagnetic Fields*. Boca Raton, FL: CRC Press.
- Pollack, H. 1975. Epidemiologic data on American personnel in the Moscow embassy. *Bull. N.Y. Acad. Med.* 55: 1182-1186.
- Polson, P. and J.H. Merritt. 1985. Cancer mortality and air force bases: a reevaluation. *J. Bioelectricity* 4: 121-127.
- Postow, E. and M.L. Swicord. 1986. Modulated fields and "window" effects. In *Handbook of Biological Effects of Electromagnetic Fields*, C. Polk and E. Postow, eds., pp. 425-460. Boca Raton, FL: CRC Press.
- Poynter, G. 1990a. The hidden hazard of traffic safety. *Law Enforcement News* 16(324): 1, 8.
- Poynter, G. 1990b. Unwitting casualties of traffic radar use. *Law Enforcement News* 16(325): 7, 10.
- Prato, F.S. et al. 1987. Attenuation of morphine-induced analgesia in mice by exposure to magnetic resonance imaging: separate effects of the static, radiofrequency and time-varying magnetic fields. *Magn. Reson. Imaging* 5: 9-14.
- Prausnitz, S. and C. Susskind. 1962. Effects of chronic microwave irradiation on mice. *IRE Trans. Bio-Med. Electron.* BME-9: 104-108.
- Preskorn, S.H., W.D. Edwards, and D.R. Justesen. 1978. Retarded tumor growth and greater longevity in mice after fetal irradiation by 2450-MHz microwaves. *J. Surg. Oncol.* 10: 483-492.
- Presman, A.S. 1970. *Electromagnetic Fields and Life*. New York: Plenum Press.
- Preston, E., E. Vavasour, and H. Assenheim. 1979. Permeability of the blood-brain barrier to mannitol in the rat following 2450 MHz microwave irradiation. *Brain Res.* 174: 109-117.
- Ragan, H. et al. 1983. Hematologic and immunologic effects of pulsed microwaves in mice. *Bioelectromagnetics* 4: 383-396.
- Rama Rao, G.R., C.A. Cain, and W.A.F. Tompkins. 1984. Effects of microwave exposure on the hamster immune system. III. Macrophage resistance to vesicular stomatitis virus infection. *Bioelectromagnetics* 5: 377-388.
- Rama Rao, G.R., C.A. Cain, and W.A.F. Tompkins. 1985. Effects of microwave exposure on the hamster immune system. IV. Spleen cell IgM hemolytic plaque formation. *Bioelectromagnetics* 6: 41-52.
- Raslear, T.G. et al. 1993. Temporal bisection in rats: the effects of high-peak-power-pulsed microwave irradiation. *Bioelectromagnetics* 14: 459-478.
- Reis, R. 1979. Potential interference with medical electronic devices. *Bull. N.Y. Acad. Med.* 55: 1216-1221.
- Remington, R.D. and M.A. Schork. 1970. *Statistics with Applications to the Biological and Health Sciences*. Englewood Cliffs, NJ: Prentice-Hall.
- Richardson, A.W., T.D. Duane, and H.M. Hines. 1948. Experimental lenticular opacities produced by microwave irradiations. *AMA Arch. Phys. Med.* 29: 765-769.
- Rigau-Perez, J.G. 1990. Summary of final discussion. *Reprod. Toxicol.* 4: 67-69.
- Roberts, N.J. 1983. Radiofrequency and microwave effects on immunological and hematopoietic systems. In *Biological Effects and Dosimetry of Nonionizing Radiation Radiofrequency and Microwave Energies*, M. Grandolfo, S. Michaelson, and A. Rindi, eds., pp. 429-459. New York: Plenum Press.
- Roberts, N.J., Lu, S.-T., and S.M. Michaelson. 1983. Human leukocyte functions and the U.S. safety standard for exposure to radiofrequency radiation. *Science* 220: 318-320.
- Roberts, N.J. and S.M. Michaelson. 1985. Epidemiological studies of human exposures to radiofrequency radiation. *Int. Arch. Occup. Environ. Health* 56: 169-178.
- Roberts, N.J., S.M. Michaelson, and S.-T. Lu. 1984. Exposure of human mononuclear leukocytes to microwave energy pulse modulated at

- 16 or 60 Hz. *IEEE Trans. Microw. Theory Tech.* MTT-32: 803-807.
- Robinette, C.D. and C. Silverman. 1977. Causes of death following occupational exposure to microwave radiation (radar) 1950-1974. In *Symposium on Biological Effects and Measurement of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D. Hazzard, ed., pp. 338-344. Rockville, MD: Bureau of Radiological Health.
- Robinette, D., C. Silverman, and S. Jablon. 1980. Effects upon health of occupational exposure to microwave radiation (radar). *Am. J. Epidemiol.* 112: 39-53.
- Rogers, S.J. 1981. Radiofrequency burn hazards in the MF/HF band. In *Proceedings of a Workshop on the Protection of Personnel Against REEM, USAFSAM Aeromedical Review* 3-81, J.C. Mitchell, ed., pp. 76-89. Cited in *Critique of the Literature on Bioeffects of Radiofrequency Radiation: A Comprehensive Review Pertinent to Air Force Operations* (Report No. USAFSAM-TR-87-3) by L. Heynick, pp. 147-148. Brooks Air Force Base, TX: U.S. Air Force School of Aerospace Medicine, 1987.
- Rohl, D. et al. 1975. The effect of radar on cardiac pacemakers. *ISA Trans.* 14: 115-117.
- Roschmann, P. 1991. Human auditory system response to pulsed radiofrequency energy in rf coils for magnetic resonance at 2.4 to 170 MHz. *Magn. Reson. Med.* 21: 197-215.
- Rose, V.E. et al. 1969. Evaluation and control of exposures in repairing microwave ovens. *Am. Ind. Hyg. Assoc. J.* 30(2): 137-142.
- Rosenthal, D.S. and S.C. Boering. 1968. Hypogonadism after microwave radiation. *JAMA* 205: 105-108.
- Rosenthal, S.W. et al. 1976. Effects of 35 and 107 GHz cw microwaves on the rabbit eye. In *Biological Effects of Electromagnetic Waves*, C.C. Johnson and M.L. Shore eds. (HEW Publication [FDA] 77-8010), pp. 110-128. Rockville, MD: Bureau of Radiological Health.
- Rotkowska, D. et al. 1993. Evaluation of the biological effects of police radar RAMER 7F. *Environ. Health Perspect.* 101: 134-136.
- Rubin, A. and W.J. Erdman. 1959. Microwave exposure of the human female pelvis during early pregnancy and prior to conception. *Am. J. Phys. Med.* 38: 219-220.
- Rugh, R. et al. 1975. Responses of the mouse to microwave radiation during estrous cycle and pregnancy. *Radiat. Res.* 62: 225-241.
- Rugh, R. 1976a. The relation of sex, age, and weight of mice to microwave radiation sensitivity. *J. Microw. Power* 11: 127-132.
- Rugh, R. 1976b. Are mouse fetuses which survive microwave radiation permanently affected thereby? *Health Phys.* 31: 33-39.
- Rugh, R. and M. McManaway. 1976. Anesthesia as an effective agent against the production of congenital anomalies in mouse fetuses exposed to electromagnetic radiation. *J. Exp. Zool.* 197: 363-368.
- Rustan, P.L., W.D. Hurt, and J.C. Mitchell. 1973. Microwave oven interference with cardiac pacemakers. *Med. Instrum.* 7: 185-188.
- Sadzikova, M.N. 1974. Clinical manifestations of reactions to microwave irradiation in various occupational groups. In *Biological Effects and Health Hazards of Microwave Radiation*, pp. 261-267. Warsaw, Poland: Polish Medical Publishers.
- Saffer, J.D. and L.A. Profano. 1992. Microwave-specific heating affects gene expression. *Bioelectromagnetics* 13: 75-78.
- Saito, M. and H.P. Schwan. 1961. The time constants of pearl-chain formation. In *Proceedings of the Fourth Annual Tri-Service Conference on the Biological Effects of Microwave Radiation*, vol. 1, M.F. Peyton, ed., pp. 85-97. New York: Plenum Press.
- Sanders, A.P. and W.T. Joines. 1983. *Effects of 200, 591, and 2450 MHz Microwaves on Cerebral Energy Metabolism* (EPA-600/S1-82-014). Springfield, VA: National Technical Information Service (PB 83-116913).
- Sanders, A.P. and W.T. Joines. 1984. The effects of hyperthermia and hyperthermia plus microwaves on rat brain energy metabolism. *Bioelectromagnetics* 5: 63-70.
- Sanders, A., W. Joines, and J. Allis. 1984. The differential effects of 200, 591, and 2450 MHz radiation on rat brain energy metabolism. *Bioelectromagnetics* 5: 419-433.
- Sanders, A., W. Joines, and J. Allis. 1985. Effects of continuous-wave, pulsed, and sinusoidal-amplitude-modulated microwaves on brain energy metabolism. *Bioelectromagnetics* 6: 89-97.
- Sanders, A., D. Schaefer, and W. Joines. 1980. Microwave effects on energy metabolism of rat brain. *Bioelectromagnetics* 1: 171-181.
- Sandström, M., B. Stenberg, and K.H. Mild. 1989. Provocation tests with ELF and VLF electromagnetic fields on patients with skin problems associated with VDT work. In *Work*

- with Display Units (Second International Scientific Conference), Montreal, Canada (Abstract 101).
- Santini, R. et al. 1988. B16 melanoma development in black mice exposed to low-level microwave radiation. *Bioelectromagnetics* 9: 105-107.
- Sastradipradja, S. and K.H. Joyner. 1990. Safety of personnel, fuels and electro-explosive devices in proximity of high power HF radar installation. In *Proceedings of Radarcon '90*, vol. 2, pp. 515-519. Adelaide, SA, Australia: Defence Sci. Technol. Organ.
- Saunders, R.D., Z.J. Sienkiewicz, and C.I. Kowalczyk. 1991a. Biological effects of electromagnetic fields and radiation. *J. Radiol. Prot.* 11: 27-42.
- Saunders, R.D., Z.J. Sienkiewicz, and C.I. Kowalczyk. 1991b. *Biological Effects of Exposure to Non-Ionising Electromagnetic Fields and Radiation* (NRPB-R240). Chilton, Oxon, UK: National Radiological Protection Board.
- Savitz, D.A. and E.E. Calle. 1987. Leukemia and occupational exposure to electromagnetic fields: review of epidemiologic studies. *J. Occup. Med.* 29: 47-51.
- Schlagel, C.J. and A. Ahmed. 1982. Evidence for genetic control of microwave-induced augmentation of complement receptor-bearing B lymphocytes. *J. Immunol.* 129: 1530-1533.
- Schnorr, T.M. 1990. NIOSH epidemiologic studies of pregnancy outcomes. *Reprod. Toxicol.* 2: 247-251.
- Schnorr, T.M. et al. 1991. Video display terminals and the risk of spontaneous abortion. *N. Engl. J. Med.* 324: 727-733.
- Schreinicke, G. et al. 1990. Stress-related changes of saliva cortisol in vdu operators. *Int. Arch. Occup. Environ. Health* 62: 319-321.
- Schrot, J., J.R. Thomas, and R.A. Banvard. 1980. Modification of the repeated acquisition of response sequences in rats by low-level microwave exposure. *Bioelectromagnetics* 1: 89-99.
- Schwan, H. 1971. Interaction of microwave and radio frequency radiation with biological systems. *IEEE Trans. Microwave Theory Tech.* MTT-19: 146-152.
- Schwan, H.P. 1982. Nonthermal cellular effects of electromagnetic fields: ac-field induced ponderomotoric forces. *Br. J. Cancer* 45(suppl. V): 220-224.
- Schwan, H.P. 1987. Biological Effects of Non-Ionizing Radiations: Cellular Properties and Interactions (Lecture No. 10). Bethesda, MD: National Council on Radiation Protection and Measurements.
- Schwan, H.P. 1992. Early history of bioelectromagnetics. *Bioelectromagnetics* 13: 453-467.
- Scialli, A.R. 1990. The history of concerns about vdt. *Reprod. Toxicol.* 4: 43-44.
- Segal, A. and R. Magin. 1982. Microwaves and the blood-brain barrier: a review. *J. Bioelectricity* 1: 351-398.
- Segal, A. and R. Magin. 1983. Letter: reply to comment on "microwaves and the blood-brain barrier." *J. Bioelectricity* 2: 89-92.
- Servantie, B. et al. 1974. Pharmacologic effects of a pulsed microwave field. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 36-45. Warsaw, Poland: Polish Medical Publishers.
- Servantie, B.M. and J. Gillard. 1983. Behavioural effects of radiofrequency energy. In *Biological Effects and Dosimetry of Nonionizing Radiation Radiofrequency and Microwave Energies*, M. Grandolfo, S.M. Michaelson, and A. Rindi, eds., pp. 393-410. New York: Plenum Press.
- Servantie, B., A.M. Servantie, and J. Etienne. 1975. Synchronization of cortical neurons by a pulsed microwave field as evidenced by spectral analysis of electrocorticograms from the white rat. *Ann. N.Y. Acad. Sci.* 247: 82-86.
- Seth, H.S. and S.M. Michaelson. 1965. Microwave cataractogenesis. *J. Occup. Med.* 9: 439-442.
- Shacklett, D.E., T.J. Tredici, D.L. Epstein. 1975. Evaluation of possible microwave-induced lens changes in the United States Air Force. *Aviat. Space Environ. Med.* 46: 1403-1406.
- Shandala, M.G., M.I. Rudnev, and I.P. Los. 1981. Investigation of biological effects of nonionizing microwave radiation. In *Proceedings of US-USSR Workshop on Physical Factors-Microwaves and Low Frequency Fields*, pp. 103-112. Research Triangle Park, NC: National Institute of Environmental Health Sciences.
- Shepich, T.J. 1990. "Safety hazard information bulletin on radiofrequency radiation-caused burns" (memorandum to regional administrators). Washington, DC: USDOL, Occupational Safety and Health Administration.
- Sheikh, K. 1986. Exposure to electromagnetic fields and the risk of leukemia. *Arch. Environ. Health* 41: 56-63.

- Shelton, W.W. and J.H. Merritt. 1981. In vitro study of microwave effects on calcium efflux in rat brain tissue. *Bioelectromagnetics* 2: 161-167.
- Shore, M.L., R.P. Felten, and A. Lamanna. 1977. The effect of repetitive prenatal exposure on development in the rat. In *Symposium on Biological Effects and Measurement of Radio Frequency/Microwaves* (HEW Publication [FDA] 77-8026), D.G. Hazzard, ed., pp. 280-289. Rockville, MD: Bureau of Radiological Health.
- Siekierzynski, M. 1974. A study of the health status of microwave workers. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 273-280. Warsaw, Poland: Polish Medical Publishers.
- Siekierzynski, M. et al. 1974. Health surveillance of personnel occupationally exposed to microwaves. III. Lens translucency. *Aerospace Med.* 45: 1146-1148.
- Sigler, A.T. et al. 1965. Radiation exposure in parents of children with mongolism (Down's Syndrome). *Bull. J. Hopkins Hosp.* 117: 374-399.
- Silverman, C. 1973. Nervous and behavioral effects of microwave radiation in humans. *Am. J. Epidemiol.* 97: 219-224.
- Silverman, C. 1980. Epidemiologic studies of microwave effects. *Proc. IEEE* 68: 78-84.
- Silverman, C. 1985. Epidemiology of microwave radiation effects in humans. In *Epidemiology and Quantitation of Environmental Risk in Humans from Radiation and Other Agents*, A. Castellani, ed., pp. 433-458. New York: Plenum Press.
- Sliney, D.H. and D.L. Conover. 1975. Nonionizing radiation. In *Industrial Environmental Health*, L.V. Cralley and P.R. Atkins, eds., pp. 157-177. New York: Academic Press.
- Smialowicz, R.J. 1987. Immunologic effects of nonionizing electromagnetic radiation. *IEEE Eng. Med. Biol. Mag.* 6: 47-51.
- Smialowicz, R.J., P.L. Brugnolotti, and M.M. Riddle. 1981. Complement receptor positive spleen cells in microwave (2450)-irradiated mice. *J. Microw. Power* 16: 73-77.
- Smialowicz, R.J. et al. 1981. Chronic exposure of rats to 100-MHz (cw) radiofrequency radiation: assessment to biological effects. *Radiat. Res.* 86: 488-505.
- Smialowicz, R.J., J.B. Kinn, and J.A. Elder. 1979. Perinatal exposure of rats to 2450-MHz cw microwave radiation: effects on lymphocytes. *Radio Sci.* 14(6S): 147-153.
- Smith, M.J. et al. 1981. An investigation of health complaints and job stress in video display operations. In *Select Research Reports on Health Issues in Video Display Terminal Operators* (NTIS PB90-103383). Springfield, VA: National Technical Information Service.
- Smith, A.B. et al. 1982. *A Preliminary Report on a Cross-Sectional Survey of VDT Users at the Baltimore Sun*. Cincinnati, OH: National Institute for Occupational Safety and Health.
- Sommer, H. and H. von Gierke. 1964. Hearing sensations in electric fields. *Aerospace Med.* 35: 834-839.
- Spalding, J.F., R.W. Freyman, and L.M. Holland. 1971. Effects of 800-MHz electromagnetic radiation on body weight, activity, hematopoiesis and life span in mice. *Health Phys.* 20: 421-424.
- Spiers, D.E. and S.C. Baummer. 1991. Thermal and metabolic responsiveness of Japanese quail embryos following periodic exposure to 2,450 MHz microwaves. *Bioelectromagnetics* 12: 225-239.
- Steel, M.C. and R.J. Sheppard. 1986. Dielectric properties of lens tissue at microwave frequencies. *Bioelectromagnetics* 7: 73-81.
- Stellman, J. and S. Stellman. 1980. Health effects of radiofrequency radiation in a cohort of physical therapists (abstract). *Am. J. Epidemiol.* 112: 442-443.
- Stewart, W. and R. Ouellet-Hellstrom. 1991. *Adverse Reproductive Events and Electromagnetic Radiation* (Johns Hopkins University #1R01 OH0 2373-01A1). Springfield, VA: National Technical Information Service (PB92-145796).
- Stewart-DeHaan, P. et al. 1983. In vitro studies of microwave-induced cataract: separation of field and heating effects. *Exp. Eye Res.* 36: 75-90.
- Stewart-DeHaan, P. et al. 1985. In vitro studies of microwave-induced cataract: reciprocity between exposure duration and dose rate for pulsed microwaves. *Exp. Eye Res.* 40: 1-13.
- Stuchly, M.A. et al. 1988. Teratological assessment of exposure to time-varying magnetic field. *Teratology* 38: 461-466.
- Sulek, K. et al. 1980. Biologic effects of microwave exposure. I. Threshold conditions for the induction of the increase in complement receptor positive (CR⁺) mouse spleen cells

- following exposure to 2450-MHz microwaves. *Radiat. Res.* 83: 127-137.
- Sullivan, D.M., O.P. Gandhi, and A. Tafflove. 1988. Use of the finite-difference time-domain method for calculating em absorption in man models. *IEEE Trans. Biomed. Eng.* 35: 179-186.
- Susser, M. 1991. What is a cause and how do we know one? a grammar for pragmatic epidemiology. *Am. J. Epidemiol.* 133: 635-648.
- Sutton, C.H. and F.B. Carroll. 1979. Effects of microwave-induced hyperthermia on the blood-brain barrier of the rat. *Radio Sci.* 14(6S): 329-334.
- Swanbeck, G. and T. Bleeker. 1989. Skin problems from visual display units. *Acta Derm. Venereol. (Stockh.)* 69: 46-51.
- Szmigielski, S. et al. 1977. In vitro and in vivo inhibition of virus multiplication by microwave hyperthermia. *Arch. Virol.* 53: 71-77.
- Szmigielski, S. et al. 1982. Accelerated development of spontaneous and benzopyrene-induced skin cancer in mice to 2450-MHz microwave radiation. *Bioelectromagnetics* 3: 179-191.
- Szmigielski, S. and J. Gil. 1989. Electromagnetic fields and neoplasms. In *Bioelectromagnetic Interaction Mechanisms, Safety Standards, Protection Guides*, G. Franceschetti, O.P. Gandhi, and M. Grandolfo, eds., pp. 81-98. New York: Plenum Press.
- Szudziński, A. et al. 1982. Acceleration of the development of benzopyrene-induced skin cancer in mice by microwave radiation. *Arch. Dermatol. Res.* 274: 303-312.
- Tafflove, A. and M.E. Brodwin. 1975. Computation of the electromagnetic fields and induced temperatures within a model of the microwave-irradiated human eye. *IEEE Trans. Microw. Theory Tech.* MTT-23: 888-896.
- Taskinen, H., P. Kyyronen, and K. Hemminki. 1990. Effects of ultrasound, shortwaves, and physical exertion on pregnancy outcome in physiotherapists. *J. Epidemiol. Community Health* 44: 196-201.
- Taylor, L.S. 1981. The mechanisms of athermal microwave biological effects. *Bioelectromagnetics* 2: 259-267.
- Teixeira-Pinto, A.A. et al. 1960. The behavior of unicellular organisms in an electromagnetic field. *Exp. Cell Res.* 20: 548-564.
- Tell, R. 1990. *An Investigation of Electric and Magnetic Fields and Operator Exposure Produced by VDTs: NIOSH VDT Epidemiology Study* (Final Report; prepared for NIOSH). Las Vegas, NV: Richard Tell Associates, Inc.
- Tengroth, B.M. 1983. Cataractogenesis induced by RF and MW energy. In *Biological Effects and Dosimetry of Nonionizing Radiation Radiofrequency and Microwave Energies*, M. Grandolfo, S. Michaelson, and A. Rindi, eds., pp. 485-500. New York: Plenum Press.
- Tengroth, B. and E. Aurell. 1974. Retinal changes in microwave workers. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 302-305. Warsaw, Poland: Polish Medical Publishers.
- Thomas, J.A. and B. Ballantyne. 1990. Occupational reproductive risks: sources, surveillance, and testing. *J. Occup. Med.* 32: 547-554.
- Thomas, J.R. et al. 1975. Effects of low-level microwave radiation on behavioral baselines. *Ann. N.Y. Acad. Sci.* 247: 425-432.
- Tikkanen, J. and O.P. Heinonen. 1991. Maternal exposure to chemical and physical factors during pregnancy and cardiovascular malformations in the offspring. *Teratology* 43: 591-600.
- Tintinalli, J.E., G. Krause, and E. Gursel. 1983. Microwave radiation injury. *Ann. Emerg. Med.* 12: 645-647.
- Tofani, S. et al. 1986. Effects of continuous low-level exposure to radiofrequency radiation on intrauterine development in rats. *Health Phys.* 51: 489-499.
- Tofani, S. et al. 1987. Reply to Lu and Michaelson regarding effects of continuous low-level exposure to radiofrequency radiation. *Health Phys.* 53: 546-547.
- Toler, J. et al. 1988. Long-term study of 435 MHz radio-frequency radiation on blood borne end points in cannulated rats part II: methods, results, and summary. *J. Microw. Power Electromag. Energy* 23: 105-136.
- Trevithick, J. et al. 1987. Histopathological studies of rabbit cornea exposed to millimeter waves. *Proceedings of the Ninth Annual Conference of the Engineering in Medicine and Biology Society* (87CH2513-0), pp. 695-697. New York: IEEE.
- Tribukait, B., E. Cekan, and L.-E. Paulsson. 1986a. Effects of pulsed magnetic fields on embryonic development in mice. In *Proceedings of the International Scientific Conference: Work with Display Units*, pp. 68-70. Stockholm, Sweden. May 12-15, 1986.
- Tribukait, B., E. Cekan, and L.-E. Paulsson. 1986b. Effects of pulsed magnetic fields on

- embryonic development in mice. *Bioelectromagnetics Society Newsletter* 66: 2-3.
- Van Ummersen, C. 1961. The effect of 2450 mc radiation on the development of the chick embryo. In *Proceedings of the Fourth Annual Tri-Service Conference on the Biological Effects of Microwave Radiation*, vol. 1, M.F. Peyton, ed., pp. 201-219. New York: Plenum Press.
- Van Ummersen, C.A. and F.C. Cogan. 1965. Experimental microwave cataracts. *Arch. Environ. Health* 11: 177-178.
- Veyret, B. et al. 1991. Antibody responses of mice exposed to low-power microwaves under combined pulse-and-amplitude modulation. *Bioelectromagnetics* 12: 47-56.
- Wangemann, R. and S. Cleary. 1976. The in vivo effects of 2.45 GHz microwave radiation on rabbit serum components and sleeping times. *Rad. Environm. Biophys.* 13: 89-103.
- Ward, T.R. et al. 1986. Brain temperature measurements in rats: a comparison of microwave and ambient temperature exposures. *Bioelectromagnetics* 7: 243-258.
- Westerholm, P. and A. Ericson. 1986. Pregnancy outcome and vdu-work in a cohort of insurance clerks. In *International Scientific Conference: Work with Display Units*, pp. 665-668. Stockholm, Sweden, May 12-15, 1986.
- Weyandt, T.B. 1992. *Evaluation of Biological and Male Reproductive Function Responses to Potential Lead Exposures in 155 MM Howitzer Crewmen*. Springfield, VA: National Technical Information Service (AD-A247 384).
- WHO. 1987. *Visual Display Terminals and Workers' Health* (WHO Offset Publication No. 99). Geneva: World Health Organization.
- WHO. 1993. *Electromagnetic Fields (300 Hz to 300 GHz)* (Environmental Health Criteria 137). Geneva: World Health Organization.
- Wike, E.L. and E.J. Martin. 1985. Comments on Frey's "data analysis reveals significant microwave-induced eye damage in humans." *J. Microw. Power* 20: 181-184.
- Wiktor-Jedrzejczak, W. et al. 1977. Immune response of mice to 2450-MHz microwave radiation: overview of immunology and empirical studies of lymphoid splenic cells. *Radio Sci.* 12(6S): 209-219.
- Wiley, M. et al. 1990. *Magnetic Field Rodent Reproductive Study* (Ontario Hydro Report #HSD-91-3). Ontario, Canada: Ontario Hydro.
- Wilkens, G.M. 1991. Nonionizing Radiation. In *Patty's Industrial Hygiene and Toxicology General Principles*, 4th ed., G.D. Clayton and F.E. Clayton, eds., pp. 657-742. New York: John Wiley and Sons.
- Williams, D.B. et al. 1955. Biologic effects studies on microwave radiation. *A.M.A. Arch. Ophthalmol.* 54: 863-874.
- Williams, R.J. and E.D. Finch. 1974. Examination of the cornea following exposure to microwave radiation. *Aerospace Med.* 45: 393-396.
- Williams, R. and T. Webb. 1980. Exposure to radio-frequency radiation from an aircraft radar unit. *Aviat. Space Environ. Med.* 51: 1243-1244.
- Williams, W. et al. 1984. Effects of 2450-MHz microwave energy on the blood-brain barrier: an overview and critique of past and present research. *IEEE Trans. Microw. Theory Tech.* MTT-32: 808-817.
- Windham, G.C. et al. 1990. Use of video display terminals during pregnancy and the risk of spontaneous abortion, low birthweight, or intrauterine growth retardation. *Am. J. Ind. Med.* 18: 675-688.
- Wright, N.A. et al. 1984. Biological studies with continuous-wave radiofrequency (28 MHz) radiation. *Radiat. Res.* 97: 468-477.
- Zaret, M.M. 1972. Clinical aspects of nonionizing radiation. *IEEE Trans. Biomed. Eng.* BME-19: 313-316.
- Zaret, M.M. 1974a. Cataracts following use of microwave oven. *N.Y. State J. Med.* 74: 2032-2034.
- Zaret, M.M. 1974b. Response by the author. *N.Y. State J. Med.* 74: 2039-2048.
- Zaret, M.M. 1975. Blindness, deafness, and vestibular dysfunction in a microwave worker. *Eye, Ear, Nose Throat Monthly* 54: 291-294.
- Zaret, M.M. 1980a. Cataracts following use of cathode ray tube displays. In *Proceedings of International Symposium of Electromagnetic Waves and Biology*, Juoy-en-Josas, France, June 30-July 4, 1980.
- Zaret, M.M. 1980b. Cataracts and visual display terminals. In *Proceedings of the Conference on Health Hazards of VDUs*, pp. 49-60. Loughborough, UK: Loughborough University of Technology, December 11, 1980.
- Zaret, M.M. 1981. Testimony before the subcommittee on investigations and oversight of the committee on science and technology of

- the U.S. House of Representatives. In *Potential Health Effects of Video Display Terminals and Radio Frequency Heaters and Sealers* (May 12, 13, 1981, No. 13), pp. 98-103. Washington, DC: U.S. Government Printing Office.
- Zaret, M.M. 1984. Cataracts and visual display units. In *Health Hazards of VDTs?*, B.G. Pearce, ed., pp. 47-59. New York: John Wiley & Sons.
- Zaret, M.M. 1991. "Electronic smog" threatens cops. *Law Enforcement News* 17(331): 6, 8.
- Zaret, M.M. and M. Eisenbud. 1961. Preliminary results of studies of the lenticular effects of microwaves among exposed personnel. In *Proceedings of the Fourth Annual Tri-Service Conference on the Biological Effects of Microwave Radiation*, vol. 1, M.F. Peyton, ed., pp. 293-308. New York: Plenum Press.
- Zaret, M.M., I.T. Kaplan, and A.M. Kay. 1969. Clinical microwave cataracts. In *Biological Effects and Health Implications of Microwave Radiation*, S.F. Cleary, ed., pp. 82-84. Washington, DC: U.S. Government Printing Office.
- Zaret, M.M. and W.Z. Snyder. 1977. Cataracts and avionic radiations. *Br. J. Ophthalmol.* 61: 380-384.
- Zydecki, S. 1974. Assessment of lens translucency in juveniles, microwave workers and age-matched groups. In *Biologic Effects and Health Hazards of Microwave Radiation*, pp. 306-308. Warsaw, Poland: Polish Medical Publishers.

E202

United States
Environmental Protection
Agency

Office of Air and Radiation &
Office of Research and
Development

402-R-95-011
March 1995



Summary and Results of the April 26-27, 1993 Radiofrequency Radiation Conference Volume 2: Papers



Summary and Results of the April 26-27, 1993 Radiofrequency Radiation Conference

Volume 2: Papers

Prepared for

Office of Air and Radiation
and Office of Research and Development
U.S. Environmental Protection Agency
401 M Street, SW
Washington, DC 20460

Under Contract

Nos. 68-D0-0102 and 68-D2-0177

DISCLAIMER

Statements, recommendations, and conclusions expressed by participants of the Radiofrequency Radiation Conference (Bethesda, Maryland, April 26 and 27, 1993) and summarized in this document are their own and do not necessarily represent the views of the U.S. Environmental Protection Agency (EPA). Furthermore, mention of trade names or commercial products does not constitute endorsement or recommendation for use by EPA.

TABLE OF CONTENTS

	Page
<i>Disclaimer</i>	ii
<i>Abstract</i>	v
 A REVIEW OF RADIOFREQUENCY ELECTRIC AND MAGNETIC FIELDS IN THE GENERAL AND WORK ENVIRONMENT: 10 kHz to 100 GHz <i>Edwin D. Mantiply et al.</i>	1
 DOSIMETRY OF RADIO FREQUENCY ELECTROMAGNETIC FIELDS <i>Arthur W. Guy</i>	25
 RF SHOCKS AND BURNS: SOME KNOWN AND UNKNOWN <i>Om P. Gandhi</i>	47
 HUMAN THERMAL RESPONSES TO RF-RADIATION INDUCED HEATING DURING MAGNETIC RESONANCE IMAGING <i>Frank G. Shellock</i>	57
 EPIDEMIOLOGIC STUDIES OF NON-IONIZING RADIOFREQUENCY EXPOSURES <i>Genevieve Matanoski</i>	77
 RESPONSES OF LABORATORY MAMMALS TO RADIOFREQUENCY RADIATION (500 kHz-100 GHz) <i>Joe A. Elder</i>	87
 RADIOFREQUENCY RADIATION EFFECTS ON CELLS <i>Stephen F. Cleary</i>	101
 ALTERATIONS IN ORNITHINE DECARBOXYLASE ACTIVITY: A CELLULAR RESPONSE TO LOW-ENERGY ELECTROMAGNETIC FIELD EXPOSURE <i>Craig V. Byus</i>	111

ABSTRACT

On April 26 and 27, 1993, the U.S. Environmental Protection Agency (EPA) Office of Air and Radiation and Office of Research and Development held a conference to assess the current knowledge of biological and human health effects of radiofrequency (RF) radiation and to address the need for and potential impact of finalization of federal guidance on human exposures to RF radiation. More than 200 people attended the conference. Attendees represented the federal government, academia, the private sector, trade associations, the media, and the public. Plenary papers presented at the meeting focused on current research findings on a variety of topics, including exposure assessment, dosimetry, biological effects, epidemiology, the basis for exposure limits, and emerging health issues. Panel discussions focused on identifying key scientific information needs for and the policy implications of the development of further EPA guidance on human exposures to RF radiation. This document, Volume 2, provides the plenary papers presented by speakers. Volume 1, under separate cover, provides a record of much of the information presented at the conference, outlines key recommendations provided to EPA by conference participants, and presents the EPA strategy for addressing RF radiation.

Two key recommendations for EPA emerged from the conference: (1) develop RF radiation exposure guidance as soon as possible, and (2) conduct additional research in a number of areas, particularly with respect to the potential for "nonthermal" effects. These recommendations were considered by EPA in its decision to proceed with the development of guidelines on human exposure to RF radiation and to develop a longer term strategy to address remaining issues. Part of this strategy has involved creating an inter-agency work group and requesting the National Council on Radiation Protection (NCRP) to assess several remaining issues. Information provided at the conference also was used as a basis for EPA comments to the Federal Communications Commission (FCC) 1993 proposal to adopt the RF radiation exposure guidelines developed in 1992 by the American National Standards Institute (ANSI) and the Institute for Electrical and Electronics Engineers (IEEE).

RF SHOCKS AND BURNS: SOME KNOWN AND UNKNOWN*

*Om P. Gandhi***

INTRODUCTION

Even though there is a great deal of data on currents induced in the human body for exposure to radiofrequency (RF) electromagnetic fields both for plane wave, relatively uniform exposures [1-5] and for nonuniform exposures in industrial settings [6-8] for free standing conditions as well as for conditions of contact with energized objects, the data is much more limited on RF shocks and burns [9-11]. This is due to the severity of the phenomenon. From anecdotal stories it is obvious that this can be a serious problem for people exposed to medium and high intensity electromagnetic (EM) fields. Following up on the pioneering work of Dalziel and colleagues [9, 10, 12, 13], Guy and others [14-17] have studied the threshold currents for perception and pain ("let-go") under conditions of continuous rather than intermittent contact with RF energized electrodes. Because of the potential for harm, currents for RF shocks and burns for conditions of intermittent contact were not a part of these studies and thus have not been examined. A limited amount of data is available for transient discharge or conditions of intermittent contact with energized conductors at 60 Hz [18,19]. From the limited data that we will present, it is likely that the stored energy levels needed for RF shocks and burns for transient discharge may diminish with increasing frequency up to 100 kHz, to values that are relatively independent of frequency for frequencies in excess of 100 kHz. Since the absence of data on startle reactions by transient discharges is one of the acknowledged weaknesses of the present day ANSI/IEEE RF safety guidelines [20], it is clear that additional experimental data is needed before this important facet of RF safety is resolved.

CONTACT HAZARDS IN THE VLF TO HF BAND (10 kHz to 100 MHz)

Ungrounded metallic objects in EM fields develop open-circuit voltages which may be written as:

$$V_{oc} = E_{inc} h_{eff} \quad (1)$$

where V_{oc} is the open circuit voltage, E_{inc} is the magnitude of the incident electric field, assumed to be relatively uniform, and h_{eff} is the effective height of the object relative to the ground. Effective height of the object is related to, but is not the same as, the physical height. Effective heights of some commonly encountered objects such as a car, van, bus, fence, metallic roof, etc. have been determined and are given in the literature [14, 17]. Effective height of a car, for example, is about 0.3 m. Effective heights are even larger for bigger or higher objects such as a school bus, metallic roof, etc. For incident electric fields of a few hundred volts per meter, open circuit voltages of tens to hundreds of volts may therefore be created between the object and the ground. Upon touching such an object, a current would flow through the human body whose magnitude will depend on the conditions of contact (contact area, grounding of the body, etc.). Chatterjee, et al. [17] did a study in which the body impedance and threshold currents needed to produce sensations of perception and pain were measured for 367 human subjects (197 males and 170 females of various age groups; 18-35, 36-50 and 51-70 years) for the frequency range 10 kHz to 3 MHz. The study included various types of contact such as finger contact (contact areas 25 and 144 mm²) and grasping a rod electrode (diameter = 1.5 cm, length = 14 cm) to simulate the holding of the door handle of a vehicle. The experimental data were used to develop graphs of average threshold incident electric fields that will cause the various sensations such as perception, let-go

* This paper was updated in October 1994.

** Department of Electrical Engineering, University of Utah, Salt Lake City, UT 84112.

(pain), or even burns for adult males and females [17, 21, 22]. Predictions were also made, based on scaling, for the corresponding threshold values for ten-year old children, since the currents for perception and pain were found to be proportional to the (contact area)^{1/4}. Taken from reference 17, a representative graph of the threshold incident electric fields for let-go (pain) for grounded adult males and ten-year old children for finger contact (contact area = 25 mm²) for various vehicles is given in Figure 1. For most of the cases the electric fields are lower than the maximum permissible exposure (MPE) limits given in the ANSI/IEEE C95.1-1992 [20] both for controlled and uncontrolled environments. Given in Figure 2 are the estimated incident electric fields for a wider area grasping contact that will result in currents needed for perception at various frequencies [17]. Average perception currents measured for 197 males for grasping contact with a cylindrical metallic rod of diameter 1.5 cm and length 14 cm (to simulate holding of a handle bar of an automobile) were used to estimate the incident electric fields that will result in such currents for grounded conditions of the subjects. Even though the RF source available at our disposal did not permit us to measure the currents for let-go under grasping contact conditions, we estimate these to be 25 to 30 percent larger than those needed for perception. This would imply that the threshold electric fields needed for let-go for grasping contact conditions would be 25 to 30 percent larger than the values given in Figure 2. From reference 17, perception is one of tingling/pricking sensation at frequencies lower than 70-100 kHz and one of warmth at the higher frequencies. Perception/let-go currents are relatively independent of frequency for frequencies higher than 100 kHz to 50-100 MHz [17, 23].

SHOCK, FIBRILLATION, AND BURNS

Shock and ventricular fibrillation have been discussed at length in the literature [10, 24-27] for 60 Hz currents. A review of the literature is given in [28]. The values for painful shock and ventricular fibrillation are presented by Dalziel for frequencies up to 10 kHz [13, 26]. It is believed that at higher frequencies, fairly large currents can pass through a human being without causing muscle or nerve stimulation [29]. These currents would, however, produce heating effects in the skin as well as damage to internal organs [25]. Becker et al. [30] and Dobbie [31] have reported values of RF currents which produce burns. Becker et al. [30] claim that 200 mA for 30 seconds produced reddening of the skin of an arm or hand of each of four human subjects, 300 mA for 20 seconds produced pain and blistering, and 400 mA for 10 seconds produced unbearable pain. In each case, the electrode was a 3.8 cm² disposable silver ECG electrode. Also, 400 mA through a 1 cm² Ferris Red Dot disposable electrode for 20 seconds produced a second-degree burn on the back of a subject's hand. This study was in reference to electrosurgery and so the frequency, though not stated in [30], is assumed to be around 500 kHz. From our results on perception and pain threshold currents beyond about 100 kHz [17], it is expected that the threshold currents for burns at other frequencies would not be any different. Dobbie [31], studying burns during surgical diathermy, reports that 100 mA through a needle electrode used in ECG monitoring over the deltoid muscle causes a unpleasantly hot sensation in 10 seconds. Becker et al. [30] report that 100 mA per square cm of skin for 10 seconds using ECG electrodes produced a burn.

Our measurements of threshold pain current on 197 male subjects indicated that an average current density of approximately 192 mA/cm² for contact with the front of the index finger for frequencies greater than or equal to 100 kHz, caused discomfort for a contact area of 25 mm². We believe that this current density would definitely have caused a burn had the subject been told to keep touching the copper plate.

It has previously been mentioned that the threshold current for perception varies with the fourth root of the contact area [17]. It is expected that the threshold current for burns will also follow the same relationship with respect to the contact area, i.e., the current density will vary as (area)^{-3/4}. For the previously studied contact areas of 144 and 25 mm², the burn thresholds can therefore be obtained by scaling 100 mA/cm² needed to cause RF burns [30], according to the relation, burn threshold current density = (area)^{-3/4}. This gives a current density of 76 mA/cm² for a contact area of 144 mm² and 283 mA/cm² for 25 mm².




Fig. 1.

Average threshold electric field for let-go (pain) for grounded adult males (solid curves) and ten-year old children (dashed curves) in finger contact with various vehicles. Contact area = 25 mm².




Fig. 2. Average threshold electric field for perception for grounded adult males (solid curves) and ten-year old children (dashed curves) in grasping contact with various vehicles.

The calculated results for threshold E-fields to cause RF burns are plotted in Figures 3 and 4 for male adults and ten-year old children for contact areas of 25 mm² and 144 mm², respectively. Since the data are not available for children, the threshold fields for children are obtained by scaling the impedance of male adults by the ratio of the height of a standard 50th percentile ten-year old child (1.38m) to the height of a standard 50th percentile male adult (1.75m) and by scaling the burn threshold current for male adults by the square of the ratio of the height of a ten-year old child to the height of a male adult.

TRANSIENT DISCHARGES

A deficiency acknowledged in the ANSI/IEEE C95.1-1992 safety guidelines [20] is that the current limits prescribed for induced and contact currents [Tables 1 and 2, parts B of ref. 20] "may not adequately protect against startle reactions caused by transient discharges when contacting an energized object". In many situations, the hazard of transient discharge may well be the most important issue for safety. Involuntary muscular reactions to transient spark discharges can cause many safety hazards. In our evaluation of construction worker safety at a U.S. Coast Guard Omega Transmitting Station (10.2-13.6 kHz), in collaboration with Robert Curtis (OSHA) and Gene Moss (NIOSH), we found that thresholds of perception, annoyance, and even spark discharges were easily exceeded for intermittent contacts with ungrounded objects (such as cables, metallic tubes, etc.) exposed to RF fields. These, of course, were a result of the energy storage (CV_{rms}^2) in these objects that could be discharged upon touching or at times even approaching these objects. Another interesting observation was that the perception threshold for finger contact for the Omega site (10.2-13.6 kHz) occurred for stored energy levels on the order of 15-20 μ J which is considerably lower than 100-150 μ J (depending on capacitance C of the objects) which is estimated at 60 Hz by experimentation with two subjects [18]. This leads us to believe that the stored energy for perception, annoyance, and spark discharges may be frequency dependent with values being considerably lower at higher frequencies than at 60 Hz where most of the data are presently available. This point of view is also shared by Dr. A. W. Guy [personal communication] who has some unpublished data at 23 kHz and feels that the perception threshold for stored energy may be even lower than 15-20 μ J that we estimate for 10.2-13.6 kHz. In another paper by Delaplace and Reilly [32], data are presented to show that threshold of spark discharge is related to energy (CV_{rms}^2) for capacitance less than 575 pF and charge (CV_{rms}^2) for $C > 575$ pF. From these limited data, most of which are at 60 Hz, it is obvious that much more data need to be obtained on the stored energy levels for transient spark discharge at various frequencies.

CONCLUDING REMARKS

Possibility of RF shocks and burns is a serious problem for personnel working close to high power transmitting antennas in the low, medium, and high frequency bands. Even though a great deal of data is available for threshold perception and let-go currents, the stored energy levels that will result in startle reactions or burns for transient discharges for intermittent contacts are not known. Some limited data points to the possibility that stored capacitive energies needed for transient discharge are likely to decrease with increasing frequencies. Since startle reactions caused by transient discharges may result in accidents, this data and instrumentation to assess which of the objects may pose such a hazard are urgently needed.